

# Effects of Different Levels of Essential Oil Mixed (Peppermint-Thyme-Anise Oil) Supplementation in the Drinking Water on the Growth Performance, Carcass Traits and Histologic Structure of Terminal Ileum in Quails <sup>[1]</sup>

Özlem KARADAĞOĞLU <sup>1</sup> ✍️ Kadir ÖNK <sup>1</sup> Tarkan ŞAHİN <sup>2</sup>  
Seyit Ali BİNGÖL <sup>3</sup> Dilek Aksu ELMALI <sup>4</sup> Özlem DURNA <sup>5</sup>

<sup>[1]</sup> This study was presented as a poster in VIIth National Animal Nutrition Congress which was held in September of 26-27, Ankara, 2013 in Turkey

<sup>1</sup> Department of Agriculture and Animal Production, Kars Vocational High School, University of Kafkas, TR-36100 Kars - TURKEY; <sup>2</sup> Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary, University of Kafkas, TR-36100 Kars - TURKEY; <sup>3</sup> Kars School of Health Sciences, Kafkas University, TR-36000 Campus, Kars - TURKEY; <sup>4</sup> Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary, University of Mustafa Kemal, TR-31040 Hatay - TURKEY; <sup>5</sup> Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary, University of Ankara, TR-06100 Ankara - TURKEY

Article Code: KVFD-2015-14390 Received: 15.09.2015 Accepted: 04.11.2015 Published Online: 04.11.2015

## Abstract

This study was conducted in order to define the effects of oregofarm (peppermint, thyme and anise oil) supplementation in the drinking water on the growth performance, carcass quality and histologic structure of terminal ileum in quails. A total of 348 Japanese quail chicks (*Coturnix coturnix japonica*) of both sexes were included in this study. They were divided into one control group and two experimental groups and each of them contained 116 Japanese quail chicks. Each group was further divided into four subgroups with 29 Japanese quail chicks. This study was finalized in six weeks. All groups were fed with basal diets and received fresh water during the experiment. The control group received non-supplemented water. The group 1 and 2 received 1.0 ml/5 L and 1.5 ml/5 L oil mixture, respectively. All experimental groups were fed with water and ad-libitum. As a result of the study, there were statistically significant differences between the feed consumption and efficiency ( $P<0.001$ ) in the end of the three weeks. Similarly, there were also statistically significant differences between same parameters ( $P<0.01$ ;  $0.05$ ) five weeks later. The body weights of quails were not significantly different from each other ( $P>0.05$ ). At the end of the study, there were statistically differences in the warm and cold carcass parameters ( $P<0.05$ ). Adding essential oil mixed were not affected on histological structure of terminal ileum ( $P>0.05$ ). Conclusively, the supplementation of oregofarm (peppermint + thyme and anise oil) has no additional effect on quail performance.

**Keywords:** Quail, Essential oil mixed, Performance, Carcass, Ileum

## İçme Suyuna Farklı Düzeylerde İlave Edilen Esansiyel Yağ Karışımının (Nane-Kekik-Anason Yağı) Bıldırcınlarda Büyüme Performansı, Karkas Parametreleri ve İleumun Histolojik Yapısı Üzerine Etkileri

### Özet

Bu araştırma, bıldırcın içme suyuna oregofarm (nane-kekik-anason yağı) ilavesinin büyüme ve besi performansı, karkas özellikleri ve ileumun histolojik yapısı üzerine etkisini belirlemek amacıyla yapılmıştır. Araştırmada her iki cinsiyette toplam 348 adet Japon bıldırcın (*Coturnix coturnix japonica*) civcivi kullanılmıştır. Her grupta 116 civciv bulunan 1 kontrol 2 deneme grubu oluşturulmuştur. Gruplar kendi aralarında 4'erli alt gruba ayrılmıştır ve her alt grup 29 adet civcivden oluşturulmuştur. Deneme altı hafta sürdürülmüştür. Bütün deneme grupları temel rasyonla beslenmiştir ve deneme boyunca temiz içme suyu sağlanmıştır. Kontrol grubunun içme suyuna oregofarm ilavesi yapılmamıştır. 1. deneme grubu ve 2. deneme grubu içme suyuna sırasıyla 1.0 mg/5 L ve 1.5 mg/5 L oregofarm ilavesi yapılmıştır. Tüm deneme gruplarına yem ve su ad-libitum olarak verilmiştir. Araştırma sonunda, yem tüketimi ve yemden yararlanma oranı bakımından 3. haftada istatistiksel açıdan önemli farklılıklar ( $P<0.001$ ) gözlenmiş olup, benzer şekilde 5. haftada da aynı değerler bakımından farklılıklar tespit edilmiştir ( $P<0.01$ ;  $0.05$ ). Denemede canlı ağırlıklar bakımından gruplar arasında istatistiksel olarak herhangi bir farklılık gözlenmemiştir ( $P>0.05$ ). Denemenin sonunda gruplar arasında sıcak ve soğuk karkas parametreleri bakımından istatistiksel olarak farklılıklar bulunmuştur ( $P<0.05$ ). Esansiyel yağ ilavesinin ileumun termal yapısı üzerine etkisi görülmemiştir ( $P>0.05$ ). Sonuç olarak, bıldırcın içme suyuna oregofarm ilavesinin performansı artırıcı ilave bir etkisinin olmadığı tespit edilmiştir.

**Anahtar sözcükler:** Bıldırcın, Esansiyel yağ karışımı, Performans, Karkas, İleum



### İletişim (Correspondence)



+90 474 2123623/125; Fax: +90 474 2239957



drozlemkaya@hotmail.com

## INTRODUCTION

For several years, the dietary use of antibiotics has been extensively used throughout the world as growth promoters in animal feeds, particularly in poultry products [1]. However, many countries have banned the use of antibiotics in animal feed as an additive since their use may lead to the antibiotic resistant bacteria which are harmful to humans [2-4]. Therefore, nutritionists and production managers have to find alternatives which have potential to alleviate the problems related to the withdrawal of antibiotics from diets and reduce enteric diseases in the poultry [5,6]. Various studies have currently focused on the research on natural feed additives in animal diets such as antioxidant [7,8], anticoccidial [9,10] or antimicrobial [11,12] plants. Recently, food additives with plant origin such as essential oils, have received considerable attention since substitutes for antibiotics are being used as a growth promoters [13]. Consequently, some plants have attracted increasing interest since they have an alternative feed additives (antimicrobial, antiparaziter, anticancer, antioxidant, antistress and appetizing) and they can replace antibiotic growth promoters [14].

Peppermint is a member of the Labiatae family and it is possibly originated from Eastern Asia. It is used widely in herbal medicine and it is believed to be beneficial particularly for the immune system. The most abundant constituent of peppermint is menthol, which has antibacterial features [15]. Peppermint is also rich source of polyphenolic compounds and hence, it could possess strong antioxidant properties [16,17]. Anise (*Pimpinella anisum* L.) is an annual aromatic plant belonging to the Apiaceae family. It has been used over the years due to its antioxidant [18], antimicrobial [19], antibacterial [20] and antifungal [21] properties. *Thymus vulgaris* L. (thyme) is also an aromatic plant belonging to Lamiaceae family. It has been a subject of considerable interest as a medicinal and therapeutic agent worldwide [22]. The pharmacological effect of thyme is attributed to carvacrol and thymol which are main bioactive components of thyme [23]. Thymol and carvacrol have considerable antimicrobial and antifungal activity [24,25]. Furthermore, they have also been reported to have antioxidant properties as well as antibacterial activity against a wide range of bacteria [26,27].

In recent years, there have been numerous studies about essential oil supplementation of dietary on poultry [28-30]. But as we know, there have not been any study supplementation of essential oil mixed in drinking water on poultry performance. The aim of the present study is to evaluate the effects of oregofarm (peppermint, thyme and anise oils) supplementation in drinking water of quails on the growth performance and carcass quality.

## MATERIAL and METHODS

This research was conducted after the approval of

Kafkas University Animal Testing Local Ethics Council (Approval Number: KAÜ-HADYEK/2012-53).

Japanese quails (*Coturnix coturnix japonica*) bred in the test unit of Kafkas University Research and Application Center. A total of 348 Japanese quail chicks were used in this study. They were divided into one control group and two experimental groups each containing 116 Japanese quail chicks. Each group was divided into four subgroups and each subgroup contained 29 Japanese quail chicks. All groups were fed *ad-libitum* with a 23% crude protein and 3080 kcal/kg metabolize energy diet. The quails received fresh water during the experiment. The control group of quails received non-supplemented water. The group 1 and 2 received 1.0 and 1.5 ml/5 L oil mixture, respectively. The active components of mixture is presented in Table 1 (Oregofarm, FARMAVET AS). It contains peppermint, thyme and anise. The composition of basal diet is presented in Table 2. The experimental period lasted in 42 days. The ratios were isocaloric and isonutrigenous. Diets were formulated to meet the NRC [31] nutrient requirements.

**Table 1.** The active components of the essential oil mixture

**Tablo 1.** Esansiyel yağ karışımının aktif bileşenleri

Components (mg/kg)	
Thymol (thyme oil)	2000
B-phellanderene (thymeoil)	1300
Limonene (thyme oil+peppermint oil)	3525
B-pinene (thyme oil+peppermint oil)	1977
Carvacrol (oreganum oil)	8910
Linalool (oreganum oil)	3645
Anethole (anise oil)	10712
Menthole (peppermint oil)	6375

**Table 2.** Composition and calculated analysis of basal diet

**Tablo 2.** Temel rasyonun bileşimi ve hesaplanan analiz değerleri

Ingredients (%)		Values Analysed (%)	
Corn	50.62	Dry Matter	90.90
Soybean meal	42.16	Crude Protein	23.02
Oil	4.17	Crude Extract	6.56
Limestone	1.34	Crude Ash	3.37
Dicalcium phosphate	1.09	Crude Fiber	2.37
Salt	0.18	ME, kcal/kg**	3080.88
Vitamin-mineral prem.*	0.20	Ca	0.95
Sodium Bicarbonate	0.08	P	0.66
DL-methionine	0.14		
L-Lysine	0.02		
Total	100		

\* Each 1 kg of Vit-min premix includes 20.000.000 IU Vit A, 3.000.000 IU Vit D<sub>3</sub>, 25 g Vitamin E, 4 g Vitamin B<sub>1</sub>, 8 g Vitamin B<sub>2</sub>, 2.5 g Vitamin B<sub>6</sub>, 20 mg Vitamin B<sub>12</sub>, 20 g Nicotinamid, 12 g Calcium-D-pantothenate, 200 g Choline Chloride, 50 g Mangane, 50 g Iron, 50 g Zinc, 10 g Copper, 0.8 g Iodine, 0.15 g Cobalt, 0.15 g Selenium; \*\* Calculated nutritional values

Crude nutrients in feed and experimental ratio were analyzed using AOAC [32] method. The levels of metabolic energy were calculated by using the formula developed by TSE [33]. At beginning (0 day) and on the 7<sup>th</sup> 14<sup>th</sup> 21<sup>st</sup>, 28<sup>th</sup>, 35<sup>th</sup> and 42<sup>nd</sup> day of the study, the body weight and body weight gain of the quails were recorded. Meanwhile, the whole feed residues of the subgroups were weighed weekly in order to measure feed consumptions and feed conversion ratios.

Eleven male and eleven female birds from each subgroups, a total of 132 chicks were randomly selected and slaughtered in order to determine their carcass quality. Then, animals were gained again and warm carcass weight was recorded. Cold carcass weight was determined upon keeping the carcasses at +4°C for 18 h. Cold carcass weight divided by the slaughter weight was calculated to determine cold carcass yield.

The ileums were taken from 6 animals of each group to examine histological. The tissues were fixed in alcohol formaldehyde then were passed series of solution which are used for routine histological process, and were embedded in paraffin. Sections taken from paraffin blocks were put on the slides and were stained with Hematoxylin Eosin (HE) and Periodic Acid Schiff (PAS) then slides were examined by light microscopy (Olympus BX-51) [34]. Length and width of villuses were measured by a micrometer and for this measurements were chosen 6 adjoining villuses at random per ileum. For villus length were taken into account between top point of ileal crypts and top of villus [35]. For width of villus was taken into account in the middle of the villus.

Statistical analyses were performed and the significance of the difference between the mean scores of the groups for body weight, body weight gain, feed consumption, feed conversion ratio, carcass weight and yield were determined by using the variance analysis method. Significant differences between the mean values of different treatments were determined by using the Duncan's multiple range tests. The statistical analysis was conducted by using the SPSS 16.0 (Inc. Chicago. IL.

USA) program. For ileum samples One-Way ANOVA in PASW statistic 18 programe was used for statistical calculations.

## RESULTS

The effects of drinking-water supplementation with peppermint+thyme and anise oils for 42 days on growth performance are shown in *Table 3*. The body weight of quail chicks was not significantly influenced by the essential oil supplementation ( $P>0.05$ ). Body weight (BW), body weight gain (BWG), feed consumption (FC) and feed conversion ratio (FCR) are shown *Table 4*. Feed consumption in control group which was not fed with essential oils was found to be higher compared to the feed consumption of experimental groups. Furthermore, the feed consumption was significantly affected by essential oils and it was found to be lower than the feed consumption of the control group ( $P<0.01$ ; 0.001). In the fifth weeks, BWGs in group II were higher when compared to other experimental groups ( $P<0.01$ ). When compared to other groups, feed conversion ratio was markedly low ( $P<0.05$ ; 0.001) in the group II (1.5 ml/5 L oregofarm) during the 3<sup>rd</sup> and 5<sup>th</sup> weeks. The overall FCR, calculated for the 6 weeks, was significantly lower than the overall FCR of other experimental groups ( $P<0.05$ ). As shown in *Table 5*, the drinking water supplementation with essential oils statistically affected the slaughter weight, warm and cold carcass ( $P<0.05$ ).

It was observed that ileum tissues were histological similar in all groups (*Fig. 1, 2, 3*). In *Table 6*, between groups, width and length of villuses were statistically determined to be similar in all groups ( $P>0.05$ ).

## DISCUSSION

According to our study, the oregofarm supplementation in the drinking water did not lead to statistically significant difference between experimental groups in terms of live body weights ( $P>0.05$ ) (*Table 3*). In terms of the results

**Table 3.** Mean body weights of the groups (g)

**Tablo 3.** Gruplarda ortalama canlı ağırlıklar (g)

Weeks	n	Control $\bar{X} \pm S_x$	n	Group I $\bar{X} \pm S_x$	n	Group II $\bar{X} \pm S_x$	P
0	116	8.41±0.08	116	8.38±0.08	116	8.38±0.07	-
1	108	14.13±0.31	107	14.07±0.34	106	13.13±0.30	-
2	100	32.43±0.94	98	33.61±0.91	100	32.30±0.89	-
3	98	65.51±1.44	95	65.78±1.48	93	65.34±1.47	-
4	97	101.33±1.92	94	100.78±1.94	91	102.98±1.87	-
5	97	140.23±2.28	94	136.71±2.37	91	142.29±2.38	-
6	96	158.88±2.48	94	155.31±2.44	91	159.27±2.77	-

-: Differences among the groups were not statistically significant ( $P>0.05$ )

**Table 4.** Mean feed consumption, weekly body weight gain and feed conversion rate\* values of groups**Tablo 4.** Gruplarda ortalama yem tüketimi, haftalık canlı ağırlık artışı ve yemden yararlanma oranı\*

Weeks	Parameters	Control X±S <sub>x</sub>	Group I X±S <sub>x</sub>	Group II X±S <sub>x</sub>	P
1	FC (g/chick)	10.01±0.23	10.12±0.45	10.19±0.18	-
	BWG (g)	5.73±0.39	5.69±0.17	4.87±0.36	-
	FCR	1.77±0.11	1.78±0.07	2.12±0.14	-
2	FC (g/chick)	48.43±1.47	49.43±3.12	44.51±3.03	-
	BWG (g)	18.19±0.56	19.14±0.56	20.39±2.01	-
	FCR	2.67±0.10	2.58±0.11	2.77±0.33	-
3	FC (g/chick)	94.59±1.46 <sup>a</sup>	93.81±1.22 <sup>a</sup>	74.12±1.64 <sup>b</sup>	***
	BWG (g)	32.76±0.40	32.13±0.70	32.16±0.71	-
	FCR	2.89±0.02 <sup>a</sup>	2.92±0.05 <sup>a</sup>	2.31±0.06 <sup>b</sup>	***
4	FC (g/chick)	120.34±1.02	117.71±4.63	112.09±1.91	-
	BWG (g)	36.46±0.83	35.22±0.94	36.27±0.45	-
	FCR	3.50±0.11	3.34±0.12	3.09±0.06	*
5	FC (g/chick)	157.51±6.10 <sup>a</sup>	141.99±1.50 <sup>b</sup>	144.17±2.32 <sup>b</sup>	**
	BWG (g)	39.27±0.24 <sup>ab</sup>	35.98±0.80 <sup>b</sup>	40.46±1.89 <sup>a</sup>	*
	FCR	4.01±0.14 <sup>a</sup>	3.95±0.08 <sup>b</sup>	3.58±0.12 <sup>b</sup>	*
6	FC (g/chick)	141.16±1.01	139.72±4.94	137.91±1.81	-
	BWG (g)	18.72 ±1.42	18.35±1.24	20.77±4.30	-
	FCR	7.80±0.66	7.72±0.57	7.32±1.10	-
1-6	FC (g/chick)	574.04±8.15 <sup>a</sup>	552.77±12.12 <sup>a</sup>	522.97±3.25 <sup>b</sup>	**
	BWG (g)	150.44±3.02	146.94±1.67	155.34±6.89	-
	FCR	3.82±0.09 <sup>a</sup>	3.76±0.05 <sup>a</sup>	3.38±0.13 <sup>b</sup>	*

a,b: Mean values on the same row followed by different letters change significantly ( $P < 0.05$ , 0.01, 0.001; respectively); -: Differences among the groups were not statistically significant ( $P > 0.05$ ); \* (kg, feed consumption/kg, body weight gain); FC: Feed Consumption, BWG: Body Weight Gain, FCR: Feed Conversion Rate

**Table 5.** Mean slaughter weight, carcass weight and carcass yields of groups**Tablo 5.** Grupların ortalama kesim ve karkas ağırlıkları ile karkas randımanları

Parameters	Control X ± S <sub>x</sub>	Group I X ± S <sub>x</sub>	Group II X ± S <sub>x</sub>	P
Slaughter weight (g)	153.89±3.68 <sup>a</sup>	141.62±3.51 <sup>b</sup>	143.75±3.90 <sup>ab</sup>	*
Warm carcass (g)	114.04±2.45 <sup>a</sup>	105.26±2.59 <sup>b</sup>	106.40±2.52 <sup>b</sup>	*
Cold carcass (g)	111.80±2.43 <sup>a</sup>	102.87±2.54 <sup>b</sup>	103.74±2.47 <sup>b</sup>	*
Warm carcass percentage (%)	74.32±0.01	74.39±0.01	74.40±0.01	-
Cold carcass percentage (%)	72.84±0.01	72.67±0.01	72.53±0.01	-


a,b: Mean values on the same row followed by different letters change significantly ( $P < 0.05$ ); -: Differences among the groups were not statistically significant ( $P > 0.05$ )

**Table 6.** The effect of different levels essential oil mixed on histologic structure of terminal ileum in Japanese quails**Tablo 6.** Japon bıldırcınlarında farklı düzeylerde esansiyel yağ karışımı ilavesinin ileumun termal yapısı üzerine etkisi

Treatment	n	Villus Length (µm)
Control ( X ± S <sub>x</sub> )	36	315.72±61.50
Group I ( X ± S <sub>x</sub> )	36	298.00±43.49
Group II ( X ± S <sub>x</sub> )	36	311.27±56.54
SEM		0.359
Treatment	n	Villus Width (µm)
Control ( X ± S <sub>x</sub> )	36	84.47±15.40
Group I ( X ± S <sub>x</sub> )	36	83.41±14.64
Group II ( X ± S <sub>x</sub> )	36	79.77±13.69
SEM		0.362


Differences among the groups were not statistically significant ( $P > 0.05$ )

of the live body weight gain, there was a statistically significant difference between the groups in the 5<sup>th</sup> week ( $P < 0.05$ ). These findings are different from the results of the study performed by Biricik et al.<sup>[13]</sup> in which they examined the effects of essential oil supplementation in the quail diets on the live body weight. However, live body weight gain results were similar to each other. In the study conducted by Parlat et al.<sup>[36]</sup> in which they searched the effects of the supplementation of virginiamycine and thyme essential oils in the diet. According to their results, the best results were obtained from quail groups which were fed with thyme essential oils. Denli et al.<sup>[37]</sup> performed a study with Japan quails and it has been shown that the thyme essential oil supplementation enhanced live body weight as well as live body weight gain according to the control group. The supplementation of the thyme + anise and rosemary extracts was shown by Tucker et al.<sup>[38]</sup>




**Fig 1.** Villuses could be observed in group 1. HE, X20. Bar 100 µm

**Şekil 1.** Grup 1'de villusların görünümü. HE, X20. Bar 100 µm



**Fig 2.** Vilusses and goblet cells could be observed in group 2. PAS, X20. Bar 100 µm

**Şekil 2.** Grup 2'de villusların ve kadeh hücrelerinin görünümü. PAS, X20. Bar 100 µm




that they increased the live body weight of broilers and decrease the death rate. Celik <sup>[39]</sup> have indicated that the different amounts of essential oil mixtures (mint + juniper + rosemary + thyme) supplemented in the diet of broilers did not affect the live body weight whereas enhanced the death rate. These results are similar to the results of our study.

In a study, different doses of anise essential oil as an alternative to antibiotics were supplemented in the diet of broilers and it has been shown that the maximum live body weight was reached when the maximum dose of

anise oil was supplemented. Therefore, they stated that anise oil can be used as a growth factor <sup>[40]</sup>. When the thyme oil was added to the diet as an alternative to antibiotics, the maximum live body weight was obtained in a group in which the thyme oil and antibiotics were supplemented together <sup>[41]</sup>. The differences between the studies can be due to the different species of animals, different usage of the extracts as well as the diets.

When the feed consumption and the rate of the feed of groups were assessed, it has been seen that there was a significant differences between groups starting from the 3<sup>rd</sup>



**Fig 3.** Cross-section of villuses, and goblet cells (arrows) could be observed in control group. PAS, X20. Bar 100 µm

**Şekil 3.** Kontrol grubu villusların enine kesiti ve kadeh hücrelerinin (oklar) görünümü. PAS, X20. Bar 100 µm

week ( $P < 0.001$ ; 0.01) (Table 4). It has been detected that the feed consumption decreased with the supplementation of the mint, thyme and anise essential oil mixture in the drinking water. Harnandez et al.<sup>[42]</sup> have reported that the different two plant extract mixtures did not have a prominent effect on the feed consumption and the rate of feed. Similarly, Tucker et al.<sup>[38]</sup> have specified that the thyme and mint oil mixtures did not have an impact on the rate of feed of the broilers. Biricik et al.<sup>[13]</sup> performed a study in which they have indicated that essential oil supplementation in quail diet did not influence statistically the feed consumption. Our results are not consistent with these findings. According to the study performed by Halle et al.<sup>[43]</sup> thyme and thyme essential oil supplementation in the diets of broilers decreased the feed consumption and the essential oil prominently enhanced the rate of feed. Our study is in agreement with the study obtained by Ghazaghi et al.<sup>[44]</sup> in which they worked with quails. Similarly, Sengün et al.<sup>[45]</sup> added thyme extract in the diet of quails and they observed the improvement in the rate of feed.

When the quails were evaluated about their carcass characteristics, there was a statistically significant difference between experimental and control groups in terms of slaughter, hot and cold carcass weights ( $P < 0.05$ ) (Table 5). According to Ghazaghi et al.<sup>[44]</sup> there was no difference between the quail groups in terms of their slaughter weights. There was no significant difference between groups related to their hot and cold carcass yields ( $P > 0.05$ ). Likewise, Biricik et al.<sup>[13]</sup> have reported that the essential oil supplementation in the diet of quails did not influence the hot and cold carcass yields. Because of these differences, due to adding different essential oil composition and added shape.

The length and the width of villus and villus goblet cell number are shown figures 1 to 3. It has been observed that there was no statistically significant difference between the experimental and the control groups related to the length and the width of the villus when the essential oil mixed are added to the drinking water of quails ( $P > 0.05$ ) (Table 6). Our results are similar to the results of Cabuk et al.<sup>[46]</sup>. In contrast to the present result, Denli et al.<sup>[37]</sup> reported that including thyme and black seed essential oil in the diet increases intestinal weight and intestinal length in quails. These differences, due to adding different essential oil composition and added shape.

Conclusively, essential oils or their mixtures have been supplemented in the diet in studies have been performed till now. In these studies, it has been aimed to examine their effects by adding them in the drinking water of animals. It has been shown that the addition of the essential oil mixtures in the drinking water of the quails decreases the feed consumption, enhances the rate of feed, and decreases the death rate. The supplementation do not have so much effect on live body weight and carcass yield whereas it has positive and prominent effects on the digestive system under the poor environmental conditions and malnutrition. Thus, further studies should be performed with different doses and under different animal husbandry systems.

## REFERENCES

1. Cabuk M, Eratak S, Alcicek A, Bozkurt M: Effects of herbal essential oil mixture as dietary supplement on egg production in quail. *Sci World J*, Article ID 573470, 2014. DOI: 10.1155/2014/573470
2. Knudsen KEB: Development of antibiotic resistance and options to replace antimicrobials in animal diets. *Proc Nutr Society*, 60, 291-299, 2001. DOI: 10.1079/PNS2001109

- 3. Hayes DJ, Jensen HH:** Lesson can be learned from Danish antibiotic ban. *Feed Stuffs*, 75 (37): 17-18, 2003.
- 4. Khaksar V, Krimpen MV, Hashemipour H, Pilevar M:** Effects of thyme essential oil on performance, some blood parameters and ileal microflora of Japanese quail. *Jpn Poult Sci Assoc*, 49, 106-110, 2012. DOI: 10.2141/jpsa.011089
- 5. Fritts CA, Waldroup PW:** Evaluation of bio-mos mannanoligosaccharide as a replacement for growth promoting antibiotics in diets for turkeys. *Int J Poult Sci*, 2, 19-22, 2003. DOI: 10.3923/ijps.2003.19.22
- 6. Ayed MH, Laamari Z, Rezik B:** Effects of incorporating an antibiotic avilamycin and a probiotic activis in broiler diets. Western Section ASAS. *Am Soc Anim Sci*, 55, 237-240, 2004.
- 7. Giannenas IA, Florou-paneri P, Botsoglou NA, Christaki E, Spais AB:** Effect of supplementing feed with oregano and/or  $\alpha$ -tocopheryl acetate on growth of broiler chickens and oxidative stability of meat. *J Anim Feed Sci*, 14, 521-535, 2005.
- 8. Florou-Paneri P, Giannenas I, Christaki E, Govaris A, Botsoglou N:** Performance of chickens and oxidative stability of the produced meat as affected by feed supplementation with oregano, vitamin C, vitamin E and their combinations. *Arch Geflugelkd*, 70(5), 232-240, 2006.
- 9. Christaki E, Florou-paneri P, Giannenas I, Papazahariadou M, Botsoglou NA, Spais AB:** Effects of mixture of herbal extracts on broiler chickens, infected with *Eimeria tenella*. *Anim Res*, 53, 137-144, 2004. DOI: 10.1051/animres:2004006
- 10. Florou-Paneri P, Christaki E, Giannenas I, Papazahariadou M, Botsoglou N.A, Spais AB:** Effect of dietary olympus tea (*Sideritis scardica*) supplementation on performance of chickens challenged with *Eimeria tenella*. *J Anim Feed Sci*, 13, 303-313, 2004.
- 11. Govaris A, Florou-Paneri P, Botsoglou E, Giannenas I, Amvrosiadis I, Botsoglou N:** The inhibitory potential of feed supplementation with rosemary and/or  $\alpha$ -tocopheryl acetate on microbial growth and lipid oxidation of turkey breast during refrigerated storage. *LWT*, 40, 331-337, 2007. DOI: 10.1016/j.lwt.2005.10.006
- 12. Botsoglou E, Govaris A, Christaki E, Botsoglou N:** Effects of dietary olive leaves and/or  $\alpha$ -tocopheryl acetate supplementation on microbial growth and lipid oxidation of turkey breast fillets during refrigerated storage. *Food Chem*, 121, 17-22, 2010. DOI: 10.1016/j.foodchem.2009.11.083
- 13. Biricik H, Yesilbag D, Gezen SS, Bulbul T:** Effects of dietary myrtle oil (*Myrtus communis* L.) supplementation on growth performance, meat oxidative stability, meat quality and erythrocyte paramaters in quails. *Revue Med Vet*, 163 (3): 134-138, 2012.
- 14. Christaki E, Bonos EM, Florou-Paneri P:** Comparative evaluation of dietary oregano, anise and olive leaves in laying Japanese guails. *Rev Bras Cienc Avic*, 13, 97-101, 2011. DOI: S1516-635X2011000200003
- 15. Schuhmacher A, Reichling J, Schnitzler P:** Virucidal effect of peppermint oil on the enveloped herpes simplex virus type 1 and type 2 *in vitro*. *Phytomed*, 10, 504-510, 2011.
- 16. Dorman HJD, Kosar K, Holm Y, Hiltunen R:** Antioxidant properties and composition of aqueous extracts from *Mentha* species, hybrids, varieties and cultivars. *J Agric Food Chem*, 51, 4563-4569, 2003. DOI: 10.1021/jf034108k
- 17. Akbari M, Torki M:** Effects of dietary chromium picolinate and peppermint essential oil on growth performance and blood biochemical parameters of broiler chicks reared under heat stress conditions. *Int J Biometeorol*, 58, 1383-1391, 2014. DOI: 10.1007/s00484-013-0740-1
- 18. Gülçin I, Oktay M, Kireççi E, Küfrecioğlu OI:** Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chem*, 83, 371-382, 2003. DOI: 10.1016/S0308-8146(03)00098-0
- 19. Al-kassie GAM:** The effect of anise and rosemary on broiler performance. *Int J Poult Sci*, 7 (3): 243-245, 2008.
- 20. Tabanca N, Bedir E, Kirimer N, Baser K.H, Khan SI, Jacob MR, Khan IA:** Antimicrobial compounds from *Pimpinella* species growing in Turkey. *Planta Med*, 69, 933-938, 2003. DOI: 10.1055/s-2003-45103
- 21. Soliman KM, Badea RI:** Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. *Food Chem Toxicol*, 40, 1669-1675, 2002. DOI: 10.1016/S0278-6915(02)00120-5
- 22. Mehdipour Z, Afsharmanesh M, Sami M:** Effects of supplemental thyme extract (*Thymus vulgaris* L.) on growth performance, intestinal populations and meat quality in Japanese quails. *Comp Clin Pathol*, 23, 1503-1508, 2014. DOI: 10.1007/s00580-013-1813-6
- 23. Grigore A, Paraschiv I, Colceru- Mihul S, Bubueanu C, Draghici E, Ichim M:** Chemical composition and antioxidant activity of *Thymus vulgaris* L. volatile oil obtained by two different methods. *Romanian Biotech Lett*, 15 (4): 5436-5443, 2010.
- 24. Twetman S, Peterson LG:** Effect of different chlorhexidine varnish regimens on mutant streptococci levels in interdental plaque and saliva. *Caries Res*, 31, 189-193, 1997. DOI: 10.1159/000262397
- 25. Basílico MZ, Basílico JC:** Inhibitory effects of some spice essential oils on *Aspergillus ochraceus* NRRL 3174 growth and ochratoxinA production. *Lett Appl Microbiol*, 29, 238-241, 1999. DOI: 10.1046/j.1365-2672.1999.00621.X
- 26. Goni P, Lopez P, Sanchez C, Gomez-Lus R, Becerril R, Nerin C:** Antimicrobial activity in the vapour phase of a combination of cinnamon and clove essential oils. *Food Chem*, 116, 982-989, 2009. DOI: 10.1016/j.foodchem.2009.03.058
- 27. Mikaili P, Mohammad Nezhady MA, Shayegh J, Asghari MH:** Study of antinociceptive effect of *Thymus vulgaris* and *Foeniculum vulgare* essential oil in mouse. *Int J Acad Res*, 2, 374-376, 2010.
- 28. Aksu T, Bozkurt AS:** Effects of dietary essential oils and/or humic acids on broiler performance, microbial population of intestinal content and antibody titres in summer season. *Kafkas Univ Vet Fak Derg*, 15, 185-190, 2009. DOI: 10.9775/kvfd.2008.89-A
- 29. Çiftçi M, Şimşek UG, Azman MA, Çerçi IH, Tonbak F:** The effects of dietary rosemary (*Rosmarinus officinalis* L.) oil supplementation on performance, carcass traits and some blood parameters of Japanese quail under heat stressed condition. *Kafkas Univ Vet Fak Derg*, 19, 595-599, 2013. DOI: 10.9775/kvfd.2012.8474
- 30. Küçükyılmaz K, Çatlı AU, Çınar M:** Etlik piliç yemlerine esansiyel yağ karışımı ilavesinin büyüme performansı, karkas randımanı ve bazı iç organ ağırlıkları üzerine etkileri. *Kafkas Univ Vet Fak Derg*, 18, 291-296, 2012. DOI: 10.9775/kvfd.2011.5443
- 31. NRC:** National Research Council, Nutrient requirements of poultry. 9<sup>th</sup> rev. ed., National Academy Press, Washington, DC 1994, pp: 44-45.
- 32. AOAC:** Official Methods of Analysis, AOAC International, Arlington, VA; 2006.
- 33. TSE:** Hayvan yemleri, metabolik (çevrilebilir enerji) tayini (Kimyasal metot). TSE No: 9610, 1991.
- 34. Luna LG:** Manual of Histologic Staining Methods of Armed Forces Institute of Pathology. 3<sup>rd</sup> ed., 222-226. Mc Graw-Hill Book Comp, New York, 1968.
- 35. Bozkurt M, Sandıkçı M:** The variations in height and width of intestinal villi and goblet cell and mitotic cell counts in chicks of different age groups. *YYU Vet Fak Derg*, 20, 5-9, 2009.
- 36. Parlat SS, Yıldız AO, Cufadar Y, Olgun O:** Effects of thyme essential oil supplementation in performance of Japanese quail exposed to experimental aflatoxicosis. *J Agr Faculty (Turkey)*, 19, 1-6, 2005.
- 37. Denli M, Okan F, Uluocak AN:** Effect of dietary supplementation of herb essential oils on the growth performance, carcass and intestinal characteristics of quail (*Coturnix coturnix japonica*). *S Afr J Anim Sci*, 34 (3): 174-179, 2004.
- 38. Tucker L:** Botanical broilers: Plant extracts enhance broiler performance. *Feed Int*, 23 (9): 26-29, 2002.
- 39. Celik R:** The effects of supplementation different levels essential oil mixed (Mint+Thyme+Thrush+Rosemary) in drinking waters on growing performance, slaughter and carcass traits in broilers. *Master Thesis*, Kafkas Univ, Health Sci Enst, 2015.
- 40. Ciftci M, Guler T, Dalkilic B, Ertas N:** The effect of anise oil (*Pimpinella Anisum* L.) on broiler performance. *Int J Poult Sci*, 4 (11): 851-855, 2005.
- 41. Simsek GU, Guler T, Cifci M, Ertas ON, Dalkilic B:** Esansiyel yağ karışımının (kekik, karanfil, anason) etlik piliçlerde canlı ağırlık, karkas ve etlerin duyuşal özellikleri üzerine etkisi. *YYU Vet Fak Derg*, 16 (2): 1-5, 2005.

- 
- 42. Hernandez F, Madrid J, Garcia V, Orengo J, Megias MD:** Influence of two plant extracts on broilers performance digestibility and digestive organ size. *Poult Sci*, 83, 169-174, 2004. DOI: 10.1093/ps/83.2.169
- 43. Halle I, Thomann R, Bauermann U, Henning M, Kohler P:** Effects of graded supplementation of herbs and essential oils in broiler feed on growth and carcass traits. *Landbauforsch Volk*, 54, 219-229, 2004
- 44. Ghazaghi M, Mehri M, Bagherzadeh-Kasmani F:** Effects of dietary mentha spicata on performance, blood metabolites, meat quality and microbial ecosystem of small intestine in growing Japanese quail. *Anim Feed Sci Technol*, 194, 89-98, 2014. DOI: 10.1016/j.anifeedsci.2014.04.014
- 45. Sengun, T, Yurtseven S, Cetin M, Kocyigit A, Sogut B:** Effect of thyme (*T. vulgaris*) extracts on fattening performance, some blood parameters, oxidative stress and DNA damage in Japanese quails. *J Anim Feed Sci*, 17, 608-620, 2008.
- 46. Cabuk, M, Eratak S, Alçiçek A, Tuğlu I:** Effect of herbal essential oil mixture on intestinal mucosal development, growth performance and weights of internal organs of quails. *J Essent Oil Bear PL*, 17, 599-606, 2014. DOI: 10.1080/0972060X.2014.935025