Kafkas Universitesi Veteriner Fakultesi Dergisi
Journal Home-Page: http://vetdergikafkas.org
E-ISSN: 1309-2251

Kafkas Univ Vet Fak Derg
32 (3): 325-335, 2026
DOI: 10.9775/kvfd.2026.36593

**Review Article**

# Targeted Prebiotics in the Control of *Eimeria* spp. to Reduce Concurrent Parasitic Zoonotic Animal Diseases in Poultry

Ahmed I. ALAJAJI [1(*)], Abdulaziz M. ALMUZAINI [1]

[1] Department of Veterinary Preventive Medicine, College of Veterinary Medicine, Qassim University, 51452 Buraidah, SAUDI ARABIA

**(*) Corresponding author:**
Ahmed I. Alajaji
Phone: +966 555138825
E-mail: ajajy@qu.edu.sa

How to cite this article?

**Alajaji AI, Almuzaini AM:** Targeted Prebiotics in the Control of *Eimeria* spp. to Reduce Concurrent Parasitic Zoonotic Animal Diseases in Poultry. *Kafkas Univ Vet Fak Derg*, 32 (3): 325-335, 2026.
DOI: 10.9775/kvfd.2026.36593

**Article ID:** KVFD-2026-36593
**Received:** 14.03.2026
**Accepted:** 13.05.2026
**Published Online:** 22.06.2026

**Abstract**

Avian coccidiosis, caused by protozoan parasites of the genus *Eimeria* remains one of the most economically devastating diseases in poultry production worldwide. Traditional methods of control are mostly based on the use of anticoccidial medication and vaccines; nevertheless, the growing development of drug-resistant strains of the parasites and governmental limitations on the usage of chemotherapeutics have initiated the development of alternative methods of control. Because of the rising concern of antimicrobial resistance, there is a need to develop alternative control strategies. There is an emergence of targeted prebiotics as promising functional feed additives that can be used to regulate the microbiota in the intestine, improve the strength of the mucosal immune system, and improve the integrity of the gut barrier. Prebiotics help to increase intestinal homeostasis and coccidial protection through selective stimulation of positive microbial populations. Moreover, because of better gut health, there could be reduced vulnerability to concomitant parasitic zoonotic pathogens, including *Toxoplasma gondii*, *Cryptosporidium parvum*, and *Giardia duodenalis*. This review investigates the application of targeted prebiotics such as fructo-oligosaccharides, mannan-oligosaccharides, inulin and β-glucans in the reduction of *Eimeria* infections, poultry performance, and in sustainable poultry production and poultry food safety.

**Keywords:** *Eimeria*, prebiotics, poultry nutrition, coccidiosis, control, gut microbiota, zoonotic parasites

## Introduction

The world poultry business has become very important in supplying the increasing demands of animal protein [1]. Poultry meat and eggs are a major source of inexpensive human foodstuffs for a developing human population [2]. Nonetheless, infectious diseases remain a major impediment to the health, production, and financial viability of poultry [3]. One of the most common and costly diseases affecting poultry in the world is avian coccidiosis, which is caused by protozoan parasites of genus *Eimeria* [4]. Coccidiosis is an intestinal infection caused by ingesting sporulated oocysts in litter, feed, or water contamination [5]. After being consumed, the parasite has several developmental stages which occur inside the epithelial cells that form the intestinal tract [6]. Some species of poultry, such as *Eimeria tenella, Eimeria maxima* and *Eimeria acervulina,* all of which attack particular parts of the intestine [7]. The consequences of infection include the destruction of epithelial cells, haemorrhage, malabsorption, a decrease in growth performance, and an enhanced susceptibility to secondary infection [8]. Coccidiosis may result in severe mortality cases, especially in young birds in severe cases [9].

The economic cost of avian coccidiosis is high and includes dwindling productivity, an upsurge in the ratios, death rates, and the expenditure on control [10]. The poultry industry has over the decades been majorly depending on anticoccidial drugs and vaccines to control the disease [11]. The chemical anticoccidials and ionophores have been extensively used in poultry feed to deter or restrict the reproductive ability of parasites [12]. Nevertheless, persistent and extensive use of such compounds has led to the development of *Eimeria* strains resistant to drugs and drug residue problems [13]. Moreover, the demand of consumers towards the antibiotic-free poultry products and the development of more stringent regulatory measures in most countries have restricted the use of chemotherapeutic agents in animal production systems [14, 15]. The challenges have prompted the need to seek alternative methods for the control of coccidiosis that are safe and sustainable [16]. The alternative control strategies include use of botanical compounds, nanoparticles, probiotics, prebiotics, peptides, immunoglobulins, vaccines, vitamins

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

and many more that can be used in avian coccidiosis and multiple other parasitic, bacterial, and viral diseases [17-19]. The effectiveness of a functional feed additive to enhance the condition of the gut and host resistance against pathogenic organisms is one such promising strategy [20,21]. The prebiotics are one of them, which have acquired significant coverage in the scientific community as they possess the capacity to selectively activate health-promoting microorganisms in the gut and enhance the function of the intestine [22].

Prebiotics can be described as non-digestible dietary factors that have health advantages to the host through their selective stimulatory effects on desirable microorganisms in the gastrointestinal tract [23]. Fructo-oligosaccharides, mannan-oligosaccharides, inulin and β-glucans are most commonly used as prebiotics in poultry [24]. These are substances that are not degraded by host enzymes but fermented by beneficial microbes along the intestine, leading to the generation of short-chain fatty acids and other metabolites, which increase the health of the gut [25]. In the intestinal microbiota, the main role is to ensure the efficiency of digestion, immune functions, and defence against infectious agents [8]. A healthy microbial environment can prevent the settlement of pathogens by competitive exclusion, generation of antimicrobial compounds, and activation of host defences [26]. On the other hand, intestinal dysbiosis may make birds susceptible to illness and undermine overall productivity [27]. *Eimeria* infection impairs intestinal integrity and changes the composition of the gut microbiota, establishing advantageous conditions in the intestine for opportunistic agents and secondary infections [28]. A promising approach to alleviate the effects of coccidiosis is the enhancement of the intestinal microbial balance by dietary interventions (e.g., specific prebiotics) [29]. Prebiotics can improve the mucosal immune system, improve the efficiency of the epithelial barrier, and slow the growth of parasites by selectively boosting the beneficial microbial populations, including *Lactobacillus* and *Bifidobacterium* [30]. Besides controlling *Eimeria* infection, there is also the possibility that better gut health reduces the chances of concomitant parasitic zoonotic infections in poultry [31].

The poultry production systems are related to various parasites of zoonotic significance: in particular, *Cryptosporidium parvum* has been identified as a major foodborne zoonotic disease [32]. They may contaminate the environment and products of poultry and threaten human well-being [33]. Dietary interventions in enhancing the intestinal defence system of poultry could lower the shedding of pathogens and contamination of the environment, and thus, result in better food safety [34]. The most recent studies have shown that specific prebiotics can tune the host immunity, alleviate gut microflora balance, and enhance resistance to illness in poultry [35]. In addition, the compounds can be used in combination with other alternative strategies, including probiotics, phytogenic additives, and vaccines to increase the overall disease control [36,37]. The recent development of interest in microbiome-based strategies is indicative of a wider adoption of sustainable and antibiotic-free systems of poultry production [38]. Although there is a growing amount of evidence supporting the effectiveness of prebiotics, but their mechanisms need to be evaluated for effective use [39]. More studies are needed to clarify the interaction of dietary ingredients and intestinal microbiota, host immune response, and the development of parasites [40]. The objective of this review is to provide a comprehensive overview of the role of targeted prebiotics in controlling *Eimeria* infections in poultry.

## Gut Health as a Central Component of Disease Resistance

Not only is the gastrointestinal tract of poultry the major place where digestive processes and the absorption occur, it is also an essential barrier against pathogenic microorganisms [41]. Healthy intestinal environment is the primary basis for keeping the overall health of poultry health, improving its growth performance, and preventing infectious diseases [8,42]. Gut health can be defined as structural integrity of intestinal epithelium, intestinal microbiota, balance and mucosal immune system efficiency [43]. The intestinal epithelium is the initial barrier to pathogenic invasion [44]. The tight junction proteins bind epithelial cells together and ensure that harmful microorganisms and toxins are not translocated into the systemic circulation [45]. The loss of intestinal integrity allows the entry of pathogens into the epithelial barrier, which causes inflammatory reactions and leads to decreased nutrient absorption rates [46]. *Eimeria* infection is a severe disorder, which leads to severe alterations in the intestinal structure by causing the death of epithelial cells throughout the intracellular parasite life cycle [47]. The effect of this damage is haemorrhage, retarded uptake of nutrients, and decreased feed efficiency [48].

Besides structural protection, the gut microbiota is critical in the protection of the host against pathogenic organisms [49]. Various microbial communities also live in the poultry intestinal tract and play a role in digestion, vitamin production, and immunity [8]. The useful micro-organisms like *Lactobacillus* and *Bifidobacterium* generate organic acids and antimicrobial substances that suppress the development of detrimental microorganisms [50]. Competitive exclusion is also a process where these microbes and pathogens compete for nutrients and

points of attachment to intestinal epithelial surfaces [51]. Another necessary element of disease resistance is the interaction of gut microbiota and host immune system [52]. The microbial metabolites in the short-chain fatty acids promote the proliferation of the intestinal immune tissues and control inflammation [53]. Normal immune responses enable the host to destroy pathogens and cause minimal damage on the tissues [54]. Nevertheless, an imbalance of microorganisms, which is also known as dysbiosis, may impair immunity and predispose to infections [55].

Avian coccidiosis has been shown to significantly alter the composition of the intestinal microbiota [56]. The lysis of the epithelial cells and cellular contents provides a good environment for opportunistic pathogens [57]. Such a microbial imbalance has the potential to worsen intestinal inflammation and increasing the severity of the disease [46]. Moreover, altered gut integrity can augment the danger of colonization by other pathogens, such as zoonotic parasites [58]. Intestinal homeostasis is thus an important aspect in proper disease management in poultry production systems [8]. Nutritional therapies to promote gut health have become more popular as an alternative to conventional antimicrobial agents [59,60]. Dietary prebiotic supplementation is one such strategy, and it has proven to be promising in tuning the gut microbiota and improve host defence programs [61].

Prebiotics facilitate the growth of beneficial microorganisms and enhance the generation of metabolites that enhance the functioning of the intestinal barrier [35]. A better microbial diversity can help to restore the intestinal balance after pathogen-induced imbalance [62]. Moreover, the better mucosal immunity enables birds to get a better response against the infection by parasites [63]. Incorporation of gut health management into poultry nutrition is one of the basic elements of sustainable disease control measures [64]. Increasing the intestinal ecosystem allows chicken farmers to decrease the occurrence and severity of such infections as coccidiosis and enhances productivity and animal welfare at the same time [13].

## Prebiotics as Functional Feed Additives

Prebiotics are non-digestible food substances that choose to stimulate the growth and metabolic activity of favourable microorganisms in the gastrointestinal tract [23]. Contrary to probiotics, which insert live microorganisms into the host, prebiotics act as food that helps to promote the growth of already existing favourable microorganisms in the gut ecosystem *(Fig. 1)* [65]. Prebiotics are being considered as a functional feed additive in the poultry nutrition field with the aim of improving the gut health, immune system, and resistance against infectious diseases [66,67]. Fructo-oligosaccharides, mannan-oligosaccharides, inulin, and β-glucans are common prebiotic compounds used in the diets of poultry [24]. These substances are not digested by the host enzymes in the upper gastrointestinal tract and appear in the lower intestine intact, where they undergo fermentation by useful microbial populations [68]. The end product of fermentation is short-chain fatty acids like acetate, propionate, and butyrate, which have significant functions in ensuring intestinal integrity as well as immunological responses [69].

The protective effects of prebiotics might include the development of microbial communities that prevent the colonization of pathogens, the reinforcement of intestinal tone, and the improvement of the immune response of the mucosa [70]. All these effects have the potential to





**Fig 1.** Targeted prebiotics in control of *Eimeria* spp. in poultry

decrease the severity of *Eimeria*-induced infections and enhance the overall performance of poultry [71].

## Fructo-Oligosaccharides

Fructo-oligosaccharides (FOS) are a short chain fructose polymers, which are usually obtained by the use of plants like chicory, onion, and garlic [72,73]. The prebiotic properties of these compounds make them a popular choice as feed additives in the poultry feed industry for the selection of beneficial intestinal microorganisms [74]. Because of their resistance to digestion by the host enzymes in the proximal gastrointestinal tract, FOS are transported to the lower intestine unfermented to be used by the microbes [68]. In chickens, FOS supplementation has been shown to stimulate the growth of good bacteria, especially *Lactobacillus* and *Bifidobacterium* [75]. *FOS fermentation* by these bacteria leads to short-chain fatty acids, of acetate, propionate, and butyrate [76]. These metabolites are important in intestinal homeostasis because they reduce intestinal pH, prevent proliferation of pathogenic microorganisms, and enhance the integrity of epithelial cells [43]. Low intestinal pH will lead to an unfriendly condition for pathogenic microbes and indirectly restrict the growth of parasites like *Eimeria* [77].

Immunomodulatory effects, which also have anticoccidial effects, are also exerted by FOS [78]. Research has shown that FOS supplementation may cause an improvement in mucosal immunity by increasing the generation of immunoglobulin A [79] and controlling the response of cytokines in the intestinal mucosa. Improved immune reactions make the recognition and killing of the stages of intracellular parasites easier [58]. Also, FOS can enhance intestinal barrier operation by inducing cell regeneration of epithelial cells and preserving tight junction integrity, which restricts the invasion of parasites and tissue injury [43].

Another important mechanism of FOS is the control of intestinal microbiota composition [80]. FOS play a role in the competitive exclusion mechanism by supporting the growth of advantageous microorganisms in the small intestine, which compete with pathogens in the intestinal environment to obtain nutrients and points of attachment, which inhibits opportunistic infections commonly related to coccidiosis [51]. Enhanced microbial balance can also decrease secondary infections which intensify the intestinal inflammation in cases of coccidiosis outbreak [81]. In combination, microbiota regulation, immune-stimulation, and increased intestinal barrier integrity FOS have the potential to be a valuable nutritional solution to prevent coccidiosis in poultry and intestinal health improvement [82].

## Mannan-Oligosaccharides

Mannan-oligosaccharides (MOS) are prebiotic compounds derived primarily from the cell walls of yeast species such as *Saccharomyces cerevisiae* [83]. These oligosaccharides consist mainly of mannose residues that possess unique biological properties beneficial for intestinal health and disease resistance in poultry [84]. One of the most important anticoccidial mechanisms of MOS involves pathogen binding and competitive exclusion [39]. Many pathogenic microorganisms possess mannose-specific lectins on their surface that enable them to attach to intestinal epithelial cells [85]. MOS molecules mimic these binding sites and act as decoy receptors, preventing pathogens from attaching to the intestinal mucosa [86]. Although *Eimeria* parasites invade epithelial cells through complex host–parasite interactions, MOS supplementation may reduce the severity of infection by limiting the colonization of secondary bacterial pathogens that exacerbate intestinal damage during coccidiosis caused by Eimeria [29]. MOS also stimulates the development of beneficial gut microbial populations, including *Lactobacillus* species [87]. Increased populations of beneficial microbes enhance fermentation processes that generate short-chain fatty acids, improving intestinal epithelial health and promoting mucosal immunity [87]. Improved intestinal microbial balance reduces inflammation and supports faster recovery from epithelial damage induced by coccidiosis [28].

Another important mechanism of MOS involves immunomodulation [88]. Research has demonstrated that MOS supplementation can stimulate both innate and adaptive immune responses in poultry [75]. Activation of macrophages, increased production of cytokines, and enhanced antibody responses contribute to improved resistance against parasitic infections [89]. Additionally, MOS may stimulate the development of gut-associated lymphoid tissues, strengthening mucosal immune defences [90]. MOS have also been associated with improvements in intestinal morphology [91]. Increased villus height and enhanced absorptive surface area contribute to improved nutrient uptake and faster recovery of intestinal tissues following coccidial damage [92]. By promoting intestinal integrity and immune competence, MOS supplementation may significantly reduce the clinical severity and production losses associated with coccidiosis in poultry [93].

The integration of prebiotics into poultry feeding programs therefore represents a promising approach for enhancing productivity while simultaneously supporting animal welfare and sustainable production practices [94].

## Inulin

Inulin is a polysaccharide that occurs naturally and is mainly composed of β-(2→1) linked fructose units with a glucose residue at its end [95]. It is often isolated out of plants like chicory root and Jerusalem artichoke and is commonly known to have prebiotic effects in

the nutrition of animals [96]. Since inulin is not digested by upper gastrointestinal tract enzymes, it will come to the large intestine where it is fermented by the good population of microbes [97]. In birds, intestinal inulin in the diet enhances the development of desirable intestinal microbes like *Bifidobacterium* and *Lactobacillus* [35]. Inulin fermentation results in the generation of short-chain fatty acids, which help to keep the intestinal pH at the optimum level and epithelial cell health [98]. One of the major products of fermentation is butyrate, which is a key source of energy to intestinal epithelial cells and is essential in the preservation of mucosal integrity [43].

It is presumed that the anticoccidial activity of inulin is triggered by a number of complementary mechanisms [12]. Inulin helps in competitive mechanisms of exclusion that prevent colonization by pathogenic microorganisms because it enhances the presence of advantageous microbial communities [51]. This microbial balance decreases gut inflammation, and it also contributes to a stable gut condition in *Eimeria*-induced infection [47].

The fermentation product of inulin improves intestinal barrier performance by increasing the growth rate of epithelial cells and strengthening tight junction proteins [98]. Inulin has been demonstrated to mediate the effects of the immune system by enhancing the activities of immune cells in the intestinal mucosa [99]. The increased production of cytokines and the improvement of secretory IgA levels augment the host's resistance to invading pathogens [100]. Besides exerting a direct effect on gut microbiota and immunity, inulin supplementation can also have an anti-oxidative stress and anti-inflammatory effect on intestinal infections [101]. Enhanced antioxidant status helps in tissue repair and healing after the damage of epithelial cells by parasitic infections [102]. These mechanisms will lead to better intestinal health, and inulin may be an effective dietary intervention during the prevention of the effects of coccidiosis in poultry production systems [5].

### β-Glucans

β-Glucans are polysaccharides made of glucose molecules connected mostly by β-(1→3) and β-(1→6) glycosidic bonds [103]. They are usually originated through yeast, fungus, cereal, and some bacteria, and have been well known in the immunomodulatory effects on animal nutrition [104]. The use of β-glucans in poultry has also been widely explored as a means of boosting immune function and resistance of the animal to infectious diseases [105]. The main action by which the β -glucans have an anticoccidial effect is through the activation of the host immune system, where the β -glucans bind to the pattern recognition receptors (DRRs) like Dectin-1 and complement receptor 3 on the immune cells like the macrophages and dendritic cells [106]. Interaction of b-glucans to these receptors causes a cascade effect of immune response that improves the host capacity to identify and exclude the invading pathogens [105]. This immune stimulation can enhance immunity against such parasites as *Eimeria* [39].

β-Glucans also provoke the synthesis of cytokines that manage inflammatory reactions and facilitating communication among the immune system cells [107]. There is better clearance of infected epithelial cells and parasite stages with increased activation of macrophages and natural killer cells in coccidiosis [39]. In addition, the β-glucans have been demonstrated to boost antibody and the growth of gut-associated lymphoid tissue [108]. The other positive impact of β-glucans is that it enhances the intestinal barrier integrity [109]. B-glucans stimulate growth of epithelial cells and facilitate processes of tissue repair that restore the intestinal architecture after being destroyed by parasitic invasion [5]. Mucosal integrity is improved, which minimizes the occurrence of secondary infections and increases the recovery of intestinal functions [44].

β-Glucans can also modify intestinal microbiota composition and make beneficial microbial populations, promoting overall gut health [110]. Immune stimulation, intestinal healing, and microbiota regulation have their combined effects that render β-glucans good nutritional interventions to curb coccidiosis in poultry production systems. MOS and β-Glucans can be used for direct immune boosting and pathogen blocking, while FOS and Inulin can be used to build a strong, parasite-resistant gut microbiome.

## Impact of Prebiotics on Poultry Performance

The addition of prebiotics to poultry diets has been largely correlated with an increase in growth performance, feed efficiency, and overall health condition [35]. It is mostly credited to improved gut health, nutrient utilization, and resistance to infectious diseases [8]. Among the most significant benefits of prebiotic supplementation, the increased feed ratio should be mentioned [111]. Prebiotics promote the growth of positive intestinal microorganisms, including *Lactobacillus,* to enhance the digestion process and the breakdown of complex nutrients into absorbable forms [112]. Prebiotic compounds are fermented to form short-chain fatty acids that trigger the growth of the intestinal epithelial and improve gut absorptive capacity [113].

Another factor that is significant in determining improved poultry performance is improved intestinal morphology [114]. Research has shown that birds fed prebiotic supplements tend to have a higher villus height and lesser depth of the crypt in the intestinal mucosa [115, 116]. These enhancement measures raise the amount of surface area used in nutrient taking leading to high growth rates and high

feed utilization [117]. Prebiotics can also work to lessen the adverse effects of intestinal pathogens on poultry production [118]. *Eimeria* infections may dramatically affect growth performance because of intestinal damage, decreased feeding levels, and decreased nutrient uptake [119]. Prebiotics can make coccidiosis milder and help the intestinal tissues to recover faster by improving intestinal microbial balance and improving immune responses [120].

Besides strengthening the resistance to coccidiosis, gut health can improve the resistance to other pathogens that affect the productivity of poultry [93]. Enhanced mucosal immunity and the balance of microbes are useful for a stable intestinal environment, preventing the colonization of pathogens [121]. Consequently, the mortality rate and the general health of the flock are significantly improved in birds that were supplemented with prebiotics *(Table 1)* [122]. Prebiotics can also lead to a higher quality carcass and safety of the product [123]. Improved intestinal health results in less shedding of pathogens and the contamination of the environment in poultry production systems [124]. This can reduce the risk of zoonotic pathogen contamination, like that of *Toxoplasma gondii* and *Cryptosporidium parvum* hence enhancing food safety and ensuring the health of the people [125,126].

## Future Prospects

The rising research in microbiome-based disease control approaches has presented novel prospects for the field of creating specific prebiotic interventions in poultry production. The current development of high-throughput sequencing tools and microbial ecology allows gaining more information about the intricate interplay of dietary ingredients, gut microbiota, host immunity, and parasitic infection. Such technologies can help identify certain groups of microbes that lead to resilience to *Eimeria*-induced infections. Future studies can involve the development of precision prebiotics that can specifically target populations of microbes that help to boost the immunity of the host

***Table 1.*** *Summary of experimental studies evaluating the effects of dietary prebiotics on poultry gut health, immune responses, growth performance, and control of coccidiosis caused by Eimeria species*

| Sr. No. | Prebiotic | Source | Poultry Type | Experimental Design | Dose / Inclusion | Duration | Target Condition | Parameters Evaluated | Mechanism | Results | Ref. |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | Mannan-oligosaccharides (MOS) | Yeast cell wall from *Saccharomyces cerevisiae* | Broiler chickens | Experimental infection with coccidia | MOS supplemented diet | 3-4 weeks | Infection with *Eimeria tenella* | Oocyst shedding, caecal lesion score | Enhanced immune response and gut protection | Reduced number of schizonts in infected birds | [127] |
| 2 | MOS + probiotic synbiotic | Yeast-derived MOS + bacterial culture | Hubbard broilers | Controlled feeding experiment (250 chicks) | Prebiotic supplementation with probiotic combination | Starter to finisher phase | Growth performance & immunity | BW gain, FCR, antibody titers, intestinal morphology | Gut microbiota modulation and immune activation | Improved bodyweight gain, villus height, and antibody levels | [128] |
| 3 | MOS and FOS | Yeast-derived MOS and plant FOS | Broiler chickens | Feeding trial (240 chicks) | 0.1-0.2% in diet | 6 weeks | Growth and meat quality | Carcass traits, lipid oxidation, cholesterol content | Improved nutrient absorption and microbial balance | Improved carcass yield and antioxidant stability | [129] |
| 4 | Fructo-oligosaccharides | Plant-derived oligosaccharides | Cobb 500 broilers | Two production cycles feeding experiment | 5 g/kg feed | 5 weeks | Immune response & microbial profile | Immune cell response, microbial counts | Lower intestinal pH and stimulation of beneficial microbes | Higher cellular immune response and reduced pathogenic bacteria | [130] |
| 5 | MOS prebiotic | Yeast cell wall polysaccharide | Commercial broilers | Controlled feeding trial | Dietary MOS supplementation | 42 days | Growth and gut histology | BW gain, FCR, gut morphology | Enhanced villus height and nutrient absorption | Improved intestinal morphology and performance | [131] |
| 6 | MOS, FOS, and inulin | Multiple prebiotic sources | Ross 308 broilers | Randomized feeding experiment | Dietary supplementation | 42 days | Metabolism & gut microbiota | Protein digestibility, gut microbial composition | Fermentation producing beneficial SCFA | Improved gut microbial diversity and nutrient metabolism | [132] |
| 7 | FOS and MOS | Plant and yeast oligosaccharides | Broiler chickens | Controlled feeding experiment | 5g/kg each | 5 weeks | Microbial balance | LAB count, pathogenic bacteria count | Competitive exclusion and acidic gut environment | Reduced *E. coli* and improved microbial balance | [133] |
| 8 | Glucomannan (prebiotic fiber) | Plant tuber (*Amorphophallus muelleri*) | Broiler chickens | Feeding experiment | 0.25% diet | Broiler growth cycle | Gut microbiota & growth performance | BW gain, FCR, microbiome sequencing | Promotion of beneficial microbes such as *Lactobacillus* spp. | Improved microbial diversity and feed efficiency | [134] |
| 9 | MOS + FOS | Prebiotic mixture | Broilers | Comparative study vs antibiotic growth promoters | 0.1-0.2% diet | 42 days | Meat quality & antioxidant status | Lipid oxidation, meat cholesterol | Improved antioxidant status and microbiota | Reduced cholesterol and oxidative damage | [129] |
| 10 | Prebiotic supplementation under stress | MOS-based prebiotic | Broiler chickens | Heat-stress experimental trial | Dietary supplementation | Several weeks | Heat stress and immune suppression | BW, mortality, WBC count | Improved gut microbiome and immune defence | Improved feed intake and reduced mortality | [122] |

and inhibit the growth of pathogenic microbes. Although these nutritional techniques can improve disease control and poultry performance, combining prebiotics with other nutritional measures like probiotics, phytogenic compounds, and vaccines could enhance the results of this effort. The alternative potential research area is the study of synbiotic preparations involving the combination of prebiotics and useful microorganisms. Such solutions can be found to have a synergistic effect to enhance intestinal health and resistance to infectious diseases. Also, long-term field experiments are required to check the efficacy of these strategies under commercial conditions of poultry production. Further investigation of microbiome-based nutrition interventions can eventually lead to poultry production systems that are sustainable and free of antibiotics.

## Conclusion

Avian coccidiosis has continued to be one of the greatest problems in poultry production, mainly because of infections by protozoan parasites of *Eimeria* genus. Conventional methods of control involving anticoccidial medication and vaccination have been effective in the management of the disease over many years but the development of drug-resistant strains of parasites and increasing limitation on the use of antimicrobial agents have led to the need to seek alternative control methods. Specific prebiotics are a novel nutritional solution to enhance intestinal condition and increase coccidiosis resistance. Fructo-oligosaccharides, mannan-oligosaccharides, inulin, and β-glucans have a positive influence through various mechanisms that include the regulation of gut microbiota, an increase in mucosal immunity, and an increase in intestinal barrier integrity. These effects lower the severity of coccidiosis but also lead to an increase in growth performance and feed efficiency in poultry. Moreover, better gut health can lead to an increased resistance to concomitant parasite zoonotic pathogens (including *Toxoplasma gondii* and *Cryptosporidium parvum*), which can improve food safety and the health of the population. Further studies on microbiome-specifically oriented nutritional interventions can offer useful options for the management of poultry diseases and the design of production systems without antibiotics.

## References

1. **Attia YA, Rahman MT, Hossain MJ, Basiouni S, Khafaga AF, Shehata AA, Hafez HM:** Poultry production and sustainability in developing countries under the COVID-19 crisis: Lessons learned. *Animals,* 12 (5): 644, 2022. DOI: 10.3390/ani12050644
2. **Istiak MS, Khaliduzzaman A:** Poultry and egg production: An overview. **In,** Khaliduzzaman A (Ed): Informatics in Poultry Production. 3-12, Springer, Singapore, 2022.
3. **Grace D, Knight-Jones TJD, Melaku A, Alders R, Jemberu WT:** The public health importance and management of infectious poultry diseases in smallholder systems in Africa. *Foods,* 13 (3):411, 2024. DOI: 10.3390/foods13030411
4. **Pal M, Rebuma T, Tolosa T:** Avian coccidiosis: A major parasitic diseas e of poultry industry. *Int J Livest Res,* 14, 1-7, 2024.
5. **Bedford MR, Apajalahti JH:** The influence of nutrition on intestinal disease with emphasis on coccidiosis. *Avian Pathol,* 51 (6): 504-520, 2022. DOI: 10.1080/03079457.2022.2098692
6. **Mesa-Pineda C, Navarro-Ruíz JL, López-Osorio S, Chaparro-Gutiérrez JJ, Gómez-Osorio LM:** Chicken coccidiosis: From the parasite lifecycle to control of the disease. *Front Vet Sci,* 8:787653, 2021. DOI: 10.3389/fvets.2021.787653
7. **Mares MM, Al-Quraishy S, Abdel-Gaber R, Murshed M:** Morphological and molecular characterization of *Eimeria* spp. infecting domestic poultry *Gallus gallus* in Riyadh City, Saudi Arabia. *Microorganisms,* 11 (3):795, 2023. DOI: 10.3390/microorganisms11030795
8. **Wickramasuriya SS, Park I, Lee K, Lee Y, Kim WH, Nam H, Lillehoj HS:** Role of physiology, immunity, microbiota, and infectious diseases in the gut health of poultry. *Vaccines,* 10 (2):172, 2022. DOI: 10.3390/vaccines10020172
9. **Ahmad R, Yu YH, Hua KF, Chen WJ, Zaborski D, Dybus A, Hsiao FSH, Cheng YH:** Management and control of coccidiosis in poultry - A review. *Anim Biosci,* 37 (1):1, 2023. DOI: 10.5713/ab.23.0189
10. **Rahmani A, Ahmed Laloui H, Kara R, Dems MA, Cherb N, Klikha A, Blake DP:** The financial cost of coccidiosis in Algerian chicken production: A major challenge for the poultry sector. *Avian Pathol,* 53 (5): 368-379, 2024. DOI: 10.1080/03079457.2024.2336091
11. **Martins RR, Silva LJG, Pereira AMPT, Esteves A, Duarte SC, Pena A:** Coccidiostats and poultry: A comprehensive review and current legislation. *Foods,* 11 (18):2738, 2022. DOI: 10.3390/foods11182738
12. **Bharti P, Tanveer S, Wani ZA, Rather SA, Parray ZA:** Insight into poultry coccidiosis: A dual focus on anticoccidial drugs and herbal plants for pharmacological and prevention strategies. *J Parasit Dis,* 49 (3): 565-582, 2025. DOI: 10.1007/s12639-025-01805-3
13. **Gao Y, Sun P, Hu D, Tang X, Zhang S, Shi F, Yan X, Yan W, Shi T, Wang S:** Advancements in understanding chicken coccidiosis: from *Eimeria* biology to innovative control strategies. *One Health Adv,* 2 (1):6, 2024. DOI: 10.1186/s44280-024-00039-x
14. **Zheng S, Li Y, Chen C, Wang N, Yang F:** Solutions to the dilemma of antibiotics use in livestock and poultry farming: Regulation policy and alternatives. *Toxics,* 13 (5):348, 2025. DOI: 10.3390/toxics13050348
15. **Magnoli AP, Parada J, Luna María J, Corti M, Escobar FM, Fernández C, Coniglio MV, Ortiz ME, Wittouck P, Watson S:** Impact of probiotic *Saccharomyces cerevisiae* var. *boulardii* RC009 alone and in combination with a phytase in broiler chickens fed with antibiotic-free diets. *Agrobiol Rec,* 16, 1-10, 2024. DOI: 10.47278/journal.abr/2024.006
16. **Zhao D, Suo J, Liang L, Liang R, Zhou R, Ding J, Liu X, Suo X, Zhang S, Tang X:** Innovative prevention and control of coccidiosis: Targeting sporogony for new control agent development. *Poult Sci,* 103 (12): 104246, 2024. DOI: 10.1016/j.psj.2024.104246
17. **Abubakar M, Oneeb M, Rashid M, Ashraf K, Chisti GA, Awan F, Sarwar NUA:** *In vitro* anthelmintic efficacy of three plant extracts against various developmental stages of *Haemonchus contortus. Pak Vet J,* 44 (2): 238-243, 2024. DOI: 10.29261/pakvetj/2024.174
18. **Bagheri E, Shori AB, Peng CW, Baba AS, Alzahrani AJ:** Phytochemical analysis and medicinal properties of some selected traditional medicinal plants. *Int J Agric Biosci,* 13 (4): 689-700, 2024. DOI: 10.47278/journal.ijab/2024.177
19. **Cissokho B, Sambe Babacar S, Diome T, Ndiaye MR, Sembene M:** *In vitro* nematocidal effects of aqueous extracts of leaves and bark of *Khaya senegalensis* and *Azadirachta indica* on the mortality of adults of *Haemonchus contortus,* cobb, 1899 (Rhabditida, Trichostrongylidae). *Agrobiol Rec,* 18, 105-112, 2024. DOI: 10.47278/journal.abr/2024.042
20. **Adedokun SA, Olojede OC:** Optimizing gastrointestinal integrity in poultry: The role of nutrients and feed additives. *Front Vet Sci,* 5:348, 2019. DOI: 10.3389/fvets.2018.00348
21. **Gul ST, Raza R, Hannan A, Khaliq S, Waheed N, Aderibigbe A:** Potential of a medicinal plant *Urtica dioica (Stinging nettle)* as a feed additive for animals and birds: A review. *Agrobiol Rec,* 2024:110-118, 2024. DOI: 10.47278/journal.abr/2024.029
22. **You S, Ma Y, Yan B, Pei W, Wu Q, Ding C, Huang C:** The promotion

mechanism of prebiotics for probiotics: A review. *Front Nutr,* 9:1000517, 2022. DOI: 10.3389/fnut.2022.1000517

23. **Chavan AR, Singh AK, Gupta RK, Nakhate SP, Poddar BJ, Gujar VV, Purohit HJ, Khardenavis AA:** Recent trends in the biotechnology of functional non-digestible oligosaccharides with prebiotic potential. *Biotechnol Genet Eng Rev,* 39 (2): 465-510, 2023. DOI: 10.1080/02648725.2022.2152627
24. **Li Z, Rafique MS, Li A, Yuan H:** Effects of probiotics on carcass, gut microbiota, and immunity of broilers. *Pak Vet J,* 45 (4): 1514-1523, 2025. DOI: 10.29261/pakvetj/2025.326
25. **Fusco W, Lorenzo MB, Cintoni M, Porcari S, Rinninella E, Kaitsas F, Lener E, Mele MC, Gasbarrini A, Collado MC:** Short-chain fatty-acid-producing bacteria: Key components of the human gut microbiota. *Nutrients,* 15 (9):2211, 2023. DOI: 10.3390/nu15092211
26. **Smith WPJ, Wucher BR, Nadell CD, Foster KR:** Bacterial defences: Mechanisms, evolution and antimicrobial resistance. *Nat Rev Microbiol,* 21 (8): 519-534, 2023. DOI: 10.1038/s41579-023-00877-3
27. **Fathima S, Shanmugasundaram R, Adams D, Selvaraj RK:** Gastrointestinal microbiota and their manipulation for improved growth and performance in chickens. *Foods,* 11 (10):1401, 2022. DOI: 10.3390/foods11101401
28. **Tomal F, Sadrin G, Gaboriaud P, Guitton E, Sedano L, Lallier N, Rossignol C, Larcher T, Rouille E, Ledevin M:** The caecal microbiota promotes the acute inflammatory response and the loss of the intestinal barrier integrity during severe *Eimeria tenella* infection. *Front Cell Infect Microbiol,* 13:1250080, 2023. DOI: 10.3389/fcimb.2023.1250080
29. **Yu H, Rahman A, Sadique MA, Batool T, Imtiaz B, Zaman MA, Riaz T, Anwar MZ, Waqas M:** Impact of *Bacillus subtilis* probiotic on growth performance, bone health, intestinal morphology, and cecal microbiota in cobb broiler chicks. *Pak Vet J,* 44 (4): 1243-1248, 2024. DOI: 10.29261/pakvetj/2024.254
30. **Al-Rashidi HS, El-Wakil ES:** Parasites and microbiota: Dual interactions and therapeutic perspectives. *Microorganisms,* 12 (10):2076, 2024. DOI: 10.3390/microorganisms12102076
31. **Mousa MR, Attia MM, Salem HM, Al-Hoshani N, Thabit H, Ibrahim MA, Albohiri HH, Khan SA, El-Saadony MT, El-Tarabily KA:** Coinfection of the gut with protozoal and metazoal parasites in broiler and laying chickens. *Poult Sci,* 103 (1):103227, 2024. DOI: 10.1016/j.psj.2023.103227
32. **Javed K, Alkheraije KA:** Cryptosporidiosis: A foodborne zoonotic disease of farm animals and humans. *Pak Vet J,* 43 (2): 213-223, 2023. DOI: 10.29261/pakvetj/2023.038
33. **Himu HA, Raihan A:** A review of the effects of intensive poultry production on the environment and human health. *Int J Vet Res,* 3 (2): 55-106, 2023.
34. **Chowdhury MAH, Ashrafudoulla M, Mevo SIU, Mizan MFR, Park SH, Ha SD:** Current and future interventions for improving poultry health and poultry food safety and security: A comprehensive review. *Compr Rev Food Sci Food Saf,* 22 (3): 1555-1596, 2023. DOI: 10.1111/1541-4337.13121
35. **Shehata AA, Yalçın S, Latorre JD, Basiouni S, Attia YA, Abd El-Wahab A, Visscher C, El-Seedi HR, Huber C, Hafez HM:** Probiotics, prebiotics, and phytogenic substances for optimizing gut health in poultry. *Microorganisms,* 10 (2):395, 2022. DOI: 10.3390/microorganisms10020395
36. **Chung ELT, Alghirani MM, Kassim NA, Ong YL, Jesse FFA, Sazili AQ, Loh TC:** Impact of *Brachiaria decumbens* leaf meal supplementation on broiler chickens raised in tropical environments in terms of growth performance, blood biochemistry, and stress biomarkers. *Braz J Poult Sci,* 26 (1): 1-8, 2024. DOI: 10.1590/1806-9061-2023-1878
37. **Fathanah SYE, Rahardja DP, Purwanti S:** Impacts of probiotic and dietary Lamtoro leaf meal on the growth performance, digestibility and small intestinal morphometry of Kampung chicken. *Int J Agric Biosci,* 13 (3): 333-339, 2024. DOI: 10.47278/journal.ijab/2024.127
38. **Leistikow KR, Beattie RE, Hristova KR:** Probiotics beyond the farm: Benefits, costs, and considerations of using antibiotic alternatives in livestock. *Front Antibiot,* 1:1003912, 2022. DOI: 10.3389/frabi.2022.1003912
39. **Lee Y, Lu M, Lillehoj HS:** Coccidiosis: Recent progress in host immunity and alternatives to antibiotic strategies. *Vaccines,* 10 (2):215, 2022. DOI: 10.3390/vaccines10020215
40. **Grondin JA, Jamal A, Mowna S, Seto T, Khan WI:** Interaction between intestinal parasites and the gut microbiota: Implications for the intestinal immune response and host defence. *Pathogens,* 13 (8): 608, 2024. DOI: 10.3390/pathogens13080608
41. **Ducatelle R, Goossens E, Eeckhaut V, Van Immerseel F:** Poultry gut health and beyond. *Anim Nutr,* 13, 240-248, 2023. DOI: 10.1016/j.aninu.2023.03.005
42. **Habib MA, Salma U, Amin MN, Rahman MG, Haque MA:** The effect of halquinol (HAL) on growth performance, carcass traits, and blood-lipid profile in broiler chickens. *Int J Agric Biosci,* 13 (1): 39-45, 2024. DOI: 10.47278/journal.ijab/2024.001
43. **Gieryńska M, Szulc-Dąbrowska L, Struzik J, Mielcarska MB, Gregorczyk-Zboroch KP:** Integrity of the intestinal barrier: The involvement of epithelial cells and microbiota-a mutual relationship. *Animals,* 12 (2):145, 2022. DOI: 10.3390/ani12020145
44. **Rogers AP, Mileto SJ, Lyras D:** Impact of enteric bacterial infections at and beyond the epithelial barrier. *Nat Rev Microbiol,* 21 (4): 260-274, 2023. DOI: 10.1038/s41579-022-00794-x
45. **Moonwiriyakit A, Pathomthongtaweechai N, Steinhagen PR, Chantawichitwong P, Satianrapapong W, Pongkorpsakol P:** Tight junctions: From molecules to gastrointestinal diseases. *Tissue Barriers,* 11 (2):2077620, 2023. DOI: 10.1080/21688370.2022.2077620
46. **Di Vincenzo F, Del Gaudio A, Petito V, Lopetuso LR, Scaldaferri F:** Gut microbiota, intestinal permeability, and systemic inflammation: A narrative review. *Intern Emerg Med,* 19 (2): 275-293, 2024. DOI: 10.1007/s11739-023-03374-w
47. **Chen P, Rehman MU, He Y, Li A, Jian F, Zhang L, Huang S:** Exploring the interplay between *Eimeria* spp. infection and the host: understanding the dynamics of gut barrier function. *Vet Q,* 45 (1): 1-22, 2025. DOI: 10.1080/01652176.2025.2452169
48. **Niu X, Ding Y, Chen S, Gooneratne R, Ju X:** Effect of immune stress on growth performance and immune functions of livestock: Mechanisms and prevention. *Animals,* 12 (7):909, 2022. DOI: 10.3390/ani12070909
49. **Wang L, Cao ZM, Zhang LL, Li JM, Lv WL:** The role of gut microbiota in some liver diseases: from an immunological perspective. *Front Immunol,* 13:923599, 2022. DOI: 10.3389/fimmu.2022.923599
50. **Zapaśnik A, Sokołowska B, Bryła M:** Role of lactic acid bacteria in food preservation and safety. *Foods,* 11 (9):1283, 2022. DOI: 10.3390/foods11091283
51. **Horrocks V, King OG, Yip AYG, Marques IM, McDonald JAK:** Role of the gut microbiota in nutrient competition and protection against intestinal pathogen colonization. *Microbiology,* 169 (8):001377, 2023. DOI: 10.1099/mic.0.001377
52. **Ahmad HI, Jabbar A, Mushtaq N, Javed Z, Hayyat MU, Bashir J, Naseeb I, Abideen ZU, Ahmad N, Chen J:** Immune tolerance vs. immune resistance: The interaction between host and pathogens in infectious diseases. *Front Vet Sci,* 9:827407, 2022. DOI: 10.3389/fvets.2022.827407
53. **Yao Y, Cai X, Fei W, Ye Y, Zhao M, Zheng C:** The role of short-chain fatty acids in immunity, inflammation and metabolism. *Critic Rev Food Sci Nutr,* 62 (1): 1-12, 2022. DOI: 10.1080/10408398.2020.1854675
54. **Cumming E, Peters C:** Immune response to infection. *Anaesth Intensive Care Med,* 25 (6): 439-443, 2024. DOI: 10.1016/j.mpaic.2024.04.003
55. **Fakharian F, Thirugnanam S, Welsh DA, Kim W-K, Rappaport J, Bittinger K, Rout N:** The role of gut dysbiosis in the loss of intestinal immune cell functions and viral pathogenesis. *Microorganisms,* 11 (7):1849, 2023. DOI: 10.3390/microorganisms11071849
56. **Jebessa E, Guo L, Chen X, Bello SF, Cai B, Girma M, Hanotte O, Nie Q:** Influence of *Eimeria maxima* coccidia infection on gut

microbiome diversity and composition of the jejunum and cecum of indigenous chicken. *Front Immunol,* 13:994224, 2022. DOI: 10.3389/fimmu.2022.994224

57. **de Hoog S, Tang C, Zhou X, Jacomel B, Lustosa B, Song Y, Kandemir H, A Ahmed S, Zhou S, Belmonte-Lopes R:** Fungal primary and opportunistic pathogens: An ecological perspective. *FEMS Microbiol Rev,* 48 (5):fuae022, 2024. DOI: 10.1093/femsre/fuae022
58. **Sardinha-Silva A, Alves-Ferreira EVC, Grigg ME:** Intestinal immune responses to commensal and pathogenic protozoa. *Front Immunol,* 13:963723, 2022. DOI: 10.3389/fimmu.2022.963723
59. **Helmy YA, Taha-Abdelaziz K, Hawwas HAE-H, Ghosh S, AlKafaas SS, Moawad MMM, Saied EM, Kassem II, Mawad AMM:** Antimicrobial resistance and recent alternatives to antibiotics for the control of bacterial pathogens with an emphasis on foodborne pathogens. *Antibiotics,* 12 (2):274, 2023. DOI: 10.3390/antibiotics12020274
60. **Azhariyanti E, Rahardja DP, Purwanti S:** Effects of early nutrition programming on post-hatching performance and small intestine characteristics of kampung chicken. *Int J Agric Biosci,* 13 (2): 128-135, 2024. DOI: 10.47278/journal.ijab/2024.103
61. **Fei Y, Chen Z, Han S, Zhang S, Zhang T, Lu Y, Berglund B, Xiao H, Li L, Yao M:** Role of prebiotics in enhancing the function of next-generation probiotics in gut microbiota. *Critic Rev Food Sci Nutr,* 63 (8): 1037-1054, 2023. DOI: 10.1080/10408398.2021.1958744
62. **Obianwuna UE, Agbai Kalu N, Wang J, Zhang H, Qi G, Qiu K, Wu S:** Recent trends on mitigative effect of probiotics on oxidative-stress-induced gut dysfunction in broilers under necrotic enteritis challenge: A review. *Antioxidants,* 12 (4):911, 2023. DOI: 10.3390/antiox12040911
63. **Sayyaf Dezfuli B, Lorenzoni M, Carosi A, Giari L, Bosi G:** Teleost innate immunity, an intricate game between immune cells and parasites of fish organs: Who wins, who loses. *Front Immunol,* 14:1250835, 2023. DOI: 10.3389/fimmu.2023.1250835
64. **Vlaicu PA, Untea AE, Varzaru I, Saracila M, Oancea AG:** Designing nutrition for health-Incorporating dietary by-products into poultry feeds to create functional foods with insights into health benefits, risks, bioactive compounds, food component functionality and safety regulations. *Foods,* 12 (21):4001, 2023. DOI: 10.3390/foods12214001
65. **Ballini A, Charitos IA, Cantore S, Topi S, Bottalico L, Santacroce L:** About functional foods: The probiotics and prebiotics state of art. *Antibiotics,* 12 (4):635, 2023. DOI: 10.3390/antibiotics12040635
66. **Amevor FK, Uyanga VA, Wu L, Xu D, Shu G, Wang Y, Zhao X:** Enhancing poultry health and productivity through the liver-gut axis with integrated nutritional and immunological approaches: a mini-review. *Front Physiol,* 16:1537099, 2025. DOI: 10.3389/fphys.2025.1537099
67. **Haryati T, Herliatika A, Sinurat AP, Wina E, Purba M, Puastuti W:** Efficacy of clove leaves, mangosteen peel extract, and liquid smoke as feed additives for native chickens. *Int J Vet Sci,* 14 (1): 107-112, 2025. DOI: 10.47278/journal.ijvs/2024.232
68. **Mukherjee A, Breselge S, Dimidi E, Marco ML, Cotter PD:** Fermented foods and gastrointestinal health: Underlying mechanisms. *Nat Rev Gastroenterol Hepatol,* 21 (4): 248-266, 2024. DOI: 10.1038/s41575-023-00869-x
69. **Liu XF, Shao JH, Liao YT, Wang LN, Jia Y, Dong PJ, Liu ZZ, He DD, Li C, Zhang X:** Regulation of short-chain fatty acids in the immune system. *Front Immunol,* 14:1186892, 2023. DOI: 10.3389/fimmu.2023.1186892
70. **Zhou P, Chen C, Patil S, Dong S:** Unveiling the therapeutic symphony of probiotics, prebiotics, and postbiotics in gut-immune harmony. *Front Nutr,* 11:1355542, 2024. DOI: 10.3389/fnut.2024.1355542
71. **Mohsin M, Zhang Z, Yin G:** Effect of probiotics on the performance and intestinal health of broiler chickens infected with *Eimeria tenella. Vaccines,* 10 (1):97, 2022. DOI: 10.3390/vaccines10010097
72. **Mudannayake DC, Jayasena DD, Wimalasiri KMS, Ranadheera CS, Ajlouni S:** Inulin fructans-food applications and alternative plant sources: A review. *Int J Food Sci Tech,* 57 (9): 5764-5780, 2022. DOI: 10.1111/ijfs.15947
73. **Abbas RZ, Raketsky V, Munir F, Mustafin B, Aubakirov M:** Botanicals in ameliorating mycotoxicosis in poultry. *Int J Vet Sci,* 13 (6): 878-888, 2024. DOI: 10.47278/journal.ijvs/2024.232
74. **Rafiq K, Tofazzal Hossain M, Ahmed R, Hasan MM, Islam R, Hossen MI, Shaha SN, Islam MR:** Role of different growth enhancers as alternative to in-feed antibiotics in poultry industry. *Front Vet Sci,* 8:794588, 2022. DOI: 10.3389/fvets.2021.794588
75. **Al-Surrayai T, Al-Khalaifah H:** Dietary supplementation of fructooligosaccharides enhanced antioxidant activity and cellular immune response in broiler chickens. *Front Vet Sci,* 9:857294, 2022. DOI: 10.3389/fvets.2022.857294
76. **Portincasa P, Bonfrate L, Vacca M, De Angelis M, Farella I, Lanza E, Khalil M, Wang DQH, Sperandio M, Di Ciaula A:** Gut microbiota and short chain fatty acids: Implications in glucose homeostasis. *Int J Mol Sci,* 23 (3):1105, 2022. DOI: 10.3390/ijms23031105
77. **Weng S, Tian E, Gao M, Zhang S, Yang G, Zhou B:** *Eimeria:* Navigating complex intestinal ecosystems. *PLoS Pathog,* 20 (11):e1012689, 2024. DOI: 10.1371/journal.ppat.1012689
78. **Costa GT, Vasconcelos QDJS, Aragão GF:** Fructooligosaccharides on inflammation, immunomodulation, oxidative stress, and gut immune response: A systematic review. *Nutr Rev,* 80 (4): 709-722, 2022. DOI: 10.1093/nutrit/nuab115
79. **Marcinkowska-Lesiak M, Motrenko M, Niewiadomski M, Głuszkiewicz I, Wojtasik-Kalinowska I, Poławska E:** Animal protein sources in Europe: Current knowledge and future perspectives - A review. *Appl Sci,* 15 (21):11749, 2025. DOI: 10.3390/app152111749
80. **Van Hul M, Cani PD:** The gut microbiota in obesity and weight management: microbes as friends or foe? *Nat Rev Endocrinol,* 19 (5): 258-271, 2023. DOI: 10.1038/s41574-022-00794-0
81. **Liu J, Guo J, Whitmore MA, Tobin I, Kim DM, Zhao Z, Zhang G:** Dynamic response of the intestinal microbiome to *Eimeria maxima*-induced coccidiosis in chickens. *Microbiol Spectr,* 12 (10): e00823-00824, 2024. DOI: 10.1128/spectrum.00823-24
82. **Prasad S, Patel B, Kumar P, Lall R:** Postbiotics: Multifunctional microbial products transforming animal health and performance. *Vet Sci,* 12 (12):1191, 2025. DOI: 10.3390/vetsci12121191
83. **Baek KR, Ramakrishnan SR, Kim SJ, Seo SO:** Yeast cell wall mannan structural features, biological activities, and production strategies. *Heliyon,* 10 (6):e27896, 2024. DOI: 10.1016/j.heliyon.2024.e27896
84. **Liu N, Shen H, Zhang F, Liu X, Xiao Q, Jiang Q, Tan B, Ma X:** Applications and prospects of functional oligosaccharides in pig nutrition: A review. *Anim Nutr,* 13, 206-215, 2023. DOI: 10.1016/j.aninu.2023.02.002
85. **Gupta A, Gupta GS:** Applications of mannose-binding lectins and mannan glycoconjugates in nanomedicine. *J Nanopart Res,* 24 (11):228, 2022. DOI: 10.1007/s11051-022-05594-1
86. **Song C, Chai Z, Chen S, Zhang H, Zhang X, Zhou Y:** Intestinal mucus components and secretion mechanisms: What we do and do not know. *Exp Mol Med,* 55 (4): 681-691, 2023. DOI: 10.1038/s12276-023-00960-y
87. **Wong CB, Huang H, Ning Y, Xiao J:** Probiotics in the new era of human milk oligosaccharides (HMOs): HMO utilization and beneficial effects of *Bifidobacterium longum* subsp. *infantis* M-63 on infant health. *Microorganisms,* 12 (5):1014, 2024. DOI: 10.3390/microorganisms12051014
88. **Strzelec M, Detka J, Mieszczak P, Sobocińska MK, Majka M:** Immunomodulation - A general review of the current state-of-the-art and new therapeutic strategies for targeting the immune system. *Front Immunol,* 14:1127704, 2023. DOI: 10.3389/fimmu.2023.1127704
89. **Obeagu EI:** Role of cytokines in immunomodulation during malaria clearance. *Ann Med Surg,* 86 (5): 2873-2882, 2024. DOI: 10.1097/ms9.0000000000002019
90. **Zhou X, Wu Y, Zhu Z, Lu C, Zhang C, Zeng L, Xie F, Zhang L, Zhou F:** Mucosal immune response in biology, disease prevention and

treatment. *Signal Transduct Target Ther,* 10 (1):7, 2025. DOI: 10.1038/s41392-024-02043-4

91. **Lu Z-Y, Feng L, Jiang W-D, Wu P, Liu Y, Jin X-W, Ren H-M, Kuang S-Y, Li S-W, Tang L:** An antioxidant supplement function exploration: Rescue of intestinal structure injury by mannan oligosaccharides after aeromonas hydrophila infection in grass carp *(Ctenopharyngodon idella). Antioxidants,* 11 (5):806, 2022. DOI: 10.3390/antiox11050806
92. **Alagbe EO, Schulze H, Adeola O:** Growth performance, nutrient digestibility, intestinal morphology, cecal mucosal cytokines and serum antioxidant responses of broiler chickens to dietary enzymatically treated yeast and coccidia challenge. *J Anim Sci Biotechnol,* 14 (1):57, 2023. DOI: 10.1186/s40104-023-00846-z
93. **El-Saadony MT, Salem HM, Attia MM, Yehia N, Abdelkader AH, Abdel-Mawgod S, Kamel NM, Alkafaas SS, Alsulami MN, Ahmed AE:** Alternatives to antibiotics against coccidiosis for poultry production: The relationship between immunity and coccidiosis management - A comprehensive review. *Ann Anim Sci,* 26 (1): 51-73, 2026. DOI: 10.2478/aoas-2025-0025
94. **Leone F, Ferrante V:** Effects of prebiotics and precision biotics on performance, animal welfare and environmental impact. A review. *Sci Total Environ,* 901:165951, 2023. DOI: 10.1016/j.scitotenv.2023.165951
95. **Tsigoriyna L, Stefanov S, Armenova N, Petrova P, Petrov K:** Microbial conversion of inulin to valuable products: The biorefinery concept. *Fermentation,* 10 (12):640, 2024. DOI: 10.3390/fermentation10120640
96. **Cornescu GM, Panaite TD, Soica C, Cismileanu A, Matache CC:** Jerusalem artichoke *(Helianthus tuberosus* L.) as a promising dietary feed ingredient for monogastric farm animals. *Appl Sci,* 13 (23):12748, 2023. DOI: 10.3390/app132312748
97. **Qin YQ, Wang LY, Yang XY, Xu YJ, Fan G, Fan YG, Ren JN, An Q, Li X:** Inulin: Properties and health benefits. *Food Func,* 14 (7): 2948-2968, 2023. DOI: 10.1039/d2fo01096h
98. **Sheng W, Ji G, Zhang L:** Immunomodulatory effects of inulin and its intestinal metabolites. *Front Immunol,* 14:1224092, 2023. DOI: 10.3389/fimmu.2023.1224092
99. **Corrêa RO, Castro PR, Fachi JL, Nirello VD, El-Sahhar S, Imada S, Pereira GV, Pral LP, Araújo NVP, Fernandes MF:** Inulin diet uncovers complex diet-microbiota-immune cell interactions remodeling the gut epithelium. *Microbiome,* 11 (1):90, 2023. DOI: 10.1186/s40168-023-01579-x
100. **Gao P, Morita N, Shinkura R:** Role of mucosal IgA antibodies as novel therapies to enhance mucosal barriers. *Semin Immunopathol*, 47 (1):1, 2025. DOI: 10.1007/s00281-024-01027-4
101. **Sahoo DK, Heilmann RM, Paital B, Patel A, Yadav VK, Wong D, Jergens AE:** Oxidative stress, hormones, and effects of natural antioxidants on intestinal inflammation in inflammatory bowel disease. *Front Endocrinol,* 14:1217165, 2023. DOI: 10.3389/fendo.2023.1217165
102. **Pawłowska M, Mila-Kierzenkowska C, Szczegielniak J, Woźniak A:** Oxidative stress in parasitic diseases - Reactive oxygen species as mediators of interactions between the host and the parasites. *Antioxidants,* 13 (1):38, 2023. DOI: 10.3390/antiox13010038
103. **Caseiro C, Dias JNR, de Andrade Fontes CMG, Bule P:** From cancer therapy to winemaking: The molecular structure and applications of β-glucans and β-1, 3-glucanases. *Int J Mol Sci,* 23 (6):3156, 2022. DOI: 10.3390/ijms23063156
104. **Gasmi A, Shanaida M, Oleshchuk O, Semenova Y, Mujawdiya PK, Ivankiv Y, Pokryshko O, Noor S, Piscopo S, Adamiv S:** Natural ingredients to improve immunity. *Pharmaceuticals,* 16 (4):528, 2023. DOI: 10.3390/ph16040528
105. **Thomas S, Rezoagli E, Abidin IZ, Major I, Murray P, Murphy EJ:** β-Glucans from yeast - Immunomodulators from novel waste resources. *Appl Sci,* 12 (10):5208, 2022. DOI: 10.3390/app12105208
106. **Muneeb M, Khan EU, Ahmad S, Naveed S, Ali M, Qazi MA, Ahmad T, Abdollahi MR:** An updated review on alternative strategies to antibiotics against necrotic enteritis in commercial broiler chickens. *Worlds Poult Sci J,* 80 (3): 821-870, 2024. DOI: 10.1080/00439339.2024.2330934
107. **Muthuramalingam K, Kim Y, Cho M:** β-glucan,"the knight of health sector": Critical insights on physiochemical heterogeneities, action mechanisms and health implications. *Critic Rev Food Sci Nutr,* 62 (25): 6908-6931, 2022. DOI: 10.1080/10408398.2021.1908221
108. **Porter D, Peggs D, McGurk C, Martin SAM:** *In-vivo* analysis of Protec™ and β-glucan supplementation on innate immune performance and intestinal health of rainbow trout. *Fish Shellfish Immunol,* 134:108573, 2023. DOI: 10.1016/j.fsi.2023.108573
109. **Karimi R, Homayoonfal M, Malekjani N, Kharazmi MS, Jafari SM:** Interaction between β-glucans and gut microbiota: A comprehensive review. *Critic Rev Food Sci Nutr,* 64 (22): 7804-7835, 2024. DOI: 10.1080/10408398.2023.2192281
110. **McCarthy C, Papada E, Kalea AZ:** The effects of cereal β-glucans on cardiovascular risk factors and the role of the gut microbiome. *Critic Rev Food Sci Nutr,* 65 (13): 2489-2505, 2025. DOI: 10.1080/10408398.2024.2345159
111. **Kiernan DP, O'Doherty JV, Sweeney T:** The effect of prebiotic supplements on the gastrointestinal microbiota and associated health parameters in pigs. *Animals,* 13 (19):3012, 2023. DOI: 10.3390/ani13193012
112. **Bedu-Ferrari C, Biscarrat P, Langella P, Cherbuy C:** Prebiotics and the human gut microbiota: From breakdown mechanisms to the impact on metabolic health. *Nutrients,* 14 (10):2096, 2022. DOI: 10.3390/nu14102096
113. **Zhang D, Jian YP, Zhang YN, Li Y, Gu LT, Sun HH, Liu MD, Zhou HL, Wang YS, Xu ZX:** Short-chain fatty acids in diseases. *Cell Commun Signal,* 21 (1):212, 2023. DOI: 10.1186/s12964-023-01219-9
114. **Xu F, Wu H, Xie J, Zeng T, Hao L, Xu W, Lu L:** The effects of fermented feed on the growth performance, antioxidant activity, immune function, intestinal digestive enzyme activity, morphology, and microflora of yellow-feather chickens. *Animals,* 13 (22):3545, 2023. DOI: 10.3390/ani13223545
115. **Rysman K, Eeckhaut V, Ducatelle R, Goossens E, Van Immerseel F:** Broiler performance correlates with gut morphology and intestinal inflammation under field conditions. *Avian Pathol,* 52 (4): 232-241, 2023. DOI: 10.1080/03079457.2023.2201169
116. **Ahmed T, Chowdhury SD, Roy BC, Das SC, Islam KMS, Ray BC, Saha S:** Ameliorative influences of prebiotic on productive performance, intestinal microbiota load, and immune response of broiler chickens. *Adv Anim Vet Sci,* 11 (5): 773-783, 2023. DOI: 10.17582/journal.aavs/2023/11.5.773.783
117. **Alharbi S, Felemban A, Abdelrahim A, Al-Dakhil M:** Agricultural and technology-based strategies to improve water-use efficiency in arid and semiarid areas. *Water,* 16 (13):1842, 2024. DOI: 10.3390/w16131842
118. **Khomayezi R, Adewole D:** Probiotics, prebiotics, and synbiotics: An overview of their delivery routes and effects on growth and health of broiler chickens. *Worlds Poult Sci J,* 78 (1): 57-81, 2022. DOI: 10.1080/00439339.2022.1988804
119. **Zhou X, Wang L, Wang Z, Zhu P, Chen Y, Yu C, Chen S, Xie Y:** Impacts of Eimeria coinfection on growth performance, intestinal health and immune responses of broiler chickens. *Vet Parasitol,* 322:110019, 2023. DOI: 10.1016/j.vetpar.2023.110019
120. **Nahed A, El-Naggar K, El-Kasrawy NI, Elblehi SS, Albadrani GM, Al-Ghadi MQ, Abdel-Daim MM:** The anticoccidial effects of probiotics and prebiotics on the live coccidia vaccine and the subsequent influence on poultry performance post-challenge with mixed Eimeria species. *Poult Sci,* 103 (12):104283, 2024. DOI: 10.1016/j.psj.2024.104283
121. **Caballero-Flores G, Pickard JM, Núñez G:** Microbiota-mediated colonization resistance: mechanisms and regulation. *Nat Rev Microbiol,* 21 (6): 347-360, 2023. DOI: 10.1038/s41579-022-00833-7
122. **Sayed Y, Hassan M, Salem HM, Al-Amry K, Eid GE:** Prophylactic influences of prebiotics on gut microbiome and immune response of heat-stressed broiler chickens. *Sci Rep,* 13 (1):13991, 2023. DOI: 10.1038/s41598-023-40997-7

123. **Lase JA, Afnan R, Wulandari Z, Estuningsih S, Sartika T, Surya, Hayanti SY, Sholikin MM, Sumantri C:** Comparison performance and carcass characteristics of native chicken breeds under free-range and intensive rearing system: A meta-analysis. *Int J Vet Sci,* 13 (6): 970-979, 2024. DOI: 10.47278/journal.ijvs/2024.192

124. **Stanley D, Bajagai YS:** Feed safety and the development of poultry intestinal microbiota. *Animals,* 12 (20):2890, 2022. DOI: 10.3390/ani12202890

125. **Almuzaini AM:** Flow of zoonotic Toxoplasmosis in food chain. *Pak Vet J,* 43 (1): 1-16, 2023. DOI: 10.29261/pakvetj/2023.010

126. **Iqbal T, Altaf S, Fatima M, Rasheed R, Laraib K, Azam M, Karamat M, Salma U, Usman S:** A narrative review on effective use of medicinal plants for the treatment of parasitic foodborne diseases. *Agrobiol Rec,* 16, 79-92, 2024. DOI: 10.47278/journal.abr/2024.016

127. **Elmusharaf MA, Bautista V, Nollet L, Beynen AC:** Effect of a mannanoligosaccharide preparation on *Eimeria tenella* infection in broiler chickens. *Int J Poult Sci,* 5 (6): 583-588, 2006. DOI: 10.3923/ijps.2006.583.588

128. **Al-Khalaifah H, Mushtaq M, Khan A, Chand N, Shah SSA, Shah M, Sultan A, Khan RU, Naz S, Abudabos A, Alhidary IA:** Effects of mannan oligosaccharide, *Bacillus clausii,* and synbiotic supplementation on growth performance, immunoglobulins, intestinal morphology, and ghrelin gene expression in broiler chickens. *Poult Sci,* 104 (11):105862, 2025. DOI: 10.1016/j.psj.2025.105862

129. **Biswas A, Mohan N, Dev K, Mir NA, Tiwari AK:** Effect of dietary mannan oligosaccharides and fructo-oligosaccharides on physico-chemical indices, antioxidant and oxidative stability of broiler chicken meat. *Sci Rep,* 11 (1):20567, 2021. DOI: 10.1038/s41598-021-99620-2

130. **Al-Khalaifa H, Al-Nasser A, Al-Surayee T, Al-Kandari S, Al-Enzi N, Al-Sharrah T, Ragheb G, Al-Qalaf S, Mohammed A:** Effect of dietary probiotics and prebiotics on the performance of broiler chickens. *Poult Sci,* 98 (10): 4465-4479, 2019. DOI: 10.3382/ps/pez282

131. **Ali A, Qureshi AS, Rehan S, Deeba F, Usman M, Ateeq MK:** Prebiotic, probiotic, and antibiotic growth promoters use in commercial broilers: A comparative study. *J Hell Vet Med Soc,* 74 (3): 5929-5936, 2023. DOI: 10.12681/jhvms.30488

132. **Nawaz A, Usman M, Ali M, Ahmad A, Jahangir M, Şahin T, Iqbal R, Khan A, Waheed S, Sarmad M:** Effects of prebiotics on metabolic activity in broilers. *Biol Clin Sci Res J,* 2024 (1):1075, 2024. DOI: 10.54112/bcsrj.v2024i1.1075

133. **Mohammed MY, Kareem KY:** A comparison study of probiotic, postbiotic and prebiotic on performance and meat quality of broilers. *Tikrit J Agric Sci,* 22 (4): 24-32, 2022. DOI: 10.25130/tjas.22.4.4

134. **Sartono HGM, Hermanto FE, Sjofjan O, Nuningtyas YF, Jatmiko YD, Alimon AR, Natsir MH:** Prebiotic effects of *Amorphophallus muelleri* on growth performance and gut microbiota in broilers. *Cogent Food Agric,* 11 (1):2566900, 2025. DOI: 10.1080/23311932.2025.2566900