

RESEARCH ARTICLE

Estimation of Protective Dose 50 of Chromatographically Purified Foot and Mouth Disease Virus Serotype A, Asia 1 and O and Immune Response of Trivalent Vaccine in Bovines

Faisal AYUB¹ , Imran ALTAF^{1(*)} , Aamir GHAFOR¹ , Ali Raza AWAN² ¹ University of Veterinary and Animal Sciences, Faculty of Veterinary Sciences, Institute of Microbiology, 54000 Lahore, PAKISTAN² University of Veterinary and Animal Sciences, Faculty of Biosciences, Institute of Biochemistry and Biotechnology, 54000 Lahore, PAKISTAN**(*) Corresponding author:**Imran Altaf
Phone: +92 3074464628
E-mail: imran.altaf@uvas.edu.pk

How to cite this article?

Ayub F, Altaf I, Ghafoor A, Awan AR: Estimation of Protective Dose 50 of Chromatographically Purified Foot and Mouth Disease Virus Serotype A, Asia 1 and O and Immune Response of Trivalent Vaccine in Bovines. *Kafkas Univ Vet Fak Derg*, 32 (3): 371-382, 2026.
DOI: 10.9775/kvfd.2026.36325**Article ID:** KVFD-2026-36325**Received:** 03.02.2026**Accepted:** 14.06.2026**Published Online:** 19.06.2026**Abstract**

Foot and Mouth Disease (FMD) is an economically significant livestock disease in developing countries such as Pakistan, where serotypes A, Asia1, and O of Foot and Mouth Disease Virus (FMDV) are prevalent. Effective disease control relies on immunization using potent vaccines, and evaluation of vaccine potency in terms of protective dose 50 (PD₅₀) is a critical step during purified vaccine formulation. The present study aimed to prepare chromatographically purified vaccines of the three prevalent serotypes in different formulations and to determine PD₅₀ values. Viruses were propagated in cell culture, biologically titrated, and chemically inactivated. To obtain highly purified antigens with reduced non-structural protein content and improved quantification compared with conventional methods, viral antigens were purified, and quantified using Size Exclusion Chromatography (SEC) with Sephacryl S-300 followed by purity check through SDS-PAGE. Five formulations of each serotype (16, 8, 4, 2, and 1 µg/dose) were prepared and inoculated into five animals per concentration. Serum samples were analyzed by SPCE to identify protected animals. PD₅₀ values were calculated for each antigen and used to formulate a trivalent vaccine. The trivalent vaccine was evaluated for structural and non-structural proteins using SPCE and 3ABC ELISA, respectively. Results demonstrated that SEC efficiently removed NSPs, as confirmed by SDS-PAGE. PD₅₀ values for serotypes A, Asia1, and O were estimated as 2.14, 1.90, and 1.63 µg/dose, respectively. Accordingly, trivalent vaccine concentrations of 6.42, 5.70, and 4.90 µg/dose (3PD₅₀) were formulated. The trivalent purified vaccine induced protective immune responses with less than 20% NSP content in SPCE and 3ABC ELISA.

Keywords: Size exclusion chromatography, Sephacryl S-300, SDS-PAGE, Nonstructural protein, PD₅₀, Vaccine

INTRODUCTION

Pakistan is an agricultural country and livestock (63.6% value addition to agro-sector) serve as the largest subsector contributing 14.97% to national gross domestic product (GDP) during year 2025-2026. More than eight million people are dependent with 35-40% income source from livestock sector. Food in the form of milk, meat & product as well skin & hides are obtained from livestock for the human consumption. It is also the mode of transportation and draught power in rural areas. There is diverse population of animals in Pakistan; buffaloes 47.7M (Million), cattle 59.7M, sheep 33.1M, goat 89.4M & camels 1.2M.

Economic growth, food security and poverty reduction are challenges of developing countries. These are due to the devastating diseases, natural disasters and climate change. Foot and Mouth Disease is one of the World Organization

of Animal Health (WOAH) listed, trans boundary & contagious disease of more than 70 species of animals. The disease is devastating not in term of only production but also economic losses due trade restrictions. Various outbreaks have been reported in neighboring countries due to uncontrolled movements of animals. Pakistan is also facing outbreaks every year in cattle population with heavy economic losses ^[1].

The disease is caused by Foot and Mouth Disease Virus (FMDV), which is positive sense of single stranded RNA member of genus *Aphthovirus* of *Picornaviridae* family. The FMDV is non-enveloped, 146S particle comprised of 8500 bases and small icosahedral protein capsid of 25-30 nm in size made up of four structural proteins each having 60 copies from VP1 to VP3 of 25-34 kDa molecular weight and VP4 of 8-10 kDa ^[2]. Virus produced ten nonstructural



proteins during replication. The FMDV is extremely labile in vitro and converted into its monomers at above 56°C and pH below 6. The FMDV have seven distinct serotypes A, O, Asia1, C, SAT-I, SAT-II and SAT-III [3-5].

Foot and mouth disease can be controlled using various strategies including the forward and backward tracing of infection origin, strict farm quarantines, restriction on movement of animals & their products, identification & culling of infected animals and vaccination of livestock herd. Mass vaccination in FMD endemic areas is key adopted measure by many countries for controlling disease. Inactivated whole culture vaccines are being used to prevent disease. The presence of nonstructural protein (NSP) of FMDV leads to dedifferentiation of vaccinated from infected animals [6,7]. Purified vaccine free from NSPs is required for FMD controlled zone as NSP are produced by infection not by the vaccination. FMD vaccines are produced using baby hamster kidney (BHK-21) cell line. While replication in the cell the FMDV translated its genome into a polyprotein i.e. co- and post-translationally sliced into 14 mature proteins out of which 04 structural protein (SP) and 10 NSPs (Lpro, 2A, 2B, 2C, 3A, 3B1-3, 3Cpro, and 3Dpol), and some cleavage intermediates [8]. Then suspension of virus including cell debris, culture media proteins and NSPs are inactivated with binary ethyleneimine (BEI) or formalin and emulsified with adjuvant to form vaccine. These proteins along with virus produce the immune response and hallmark of differentiation between vaccinated and infected animals. As per WOAHP requirement vaccine must not produce any anti-NSP antibodies after repeated immunization [9].

A potent vaccine formulation depends upon purity and quantity of antigen used. The immunogenicity of FMD vaccines is primarily associated with the presence of intact and purified 146S viral particles [10]. Therefore, concentration and purification of harvested virus are essential steps for efficient storage, quantification of antigen and development of potent vaccines. Various methods, such as precipitation of antigen using polyethylene glycol (PEG) and ammonium sulfate, and ultrafiltration through membrane with particular molecular weight cut off (MWCO) capacity, have been employed for the antigen concentration [11]. For purification and quantification of FMD virus, sucrose density gradient (SDG) ultracentrifugation has traditionally been reference method. This technique separate biomolecules on the basis of size and sedimentation characteristics using linear sucrose gradient approaches [12]. However, SDG ultracentrifugation require specialized equipment and highly skilled, involve lengthy process of 64 h, has high operating and maintenance costs, and has a low sample volume capacity, making it difficult to adopt for vaccine production on an industrial scale [13].

To overcome these limitations Size exclusion chromatography (SEC) has emerged as an alternative approach for purification and quantification of biomolecules [14]. Compared with SDG ultracentrifugation, SEC is a simpler, more economical, and less time-consuming technique that can be readily scaled up by increasing the sample-to-column volume ratio [15]. Furthermore SEC has been successfully utilized for both purification and quantification of FMDV, and has demonstrated a significant correlation value of 0.9669 with SDG-based measurement [16]. The technique utilizes a column packed with porous resin beads where smaller molecules enter pores and elute later, while larger molecules excluded from the pores and elute earlier. These characteristics make SEC a practical and promising method for purification and quantification of FMDV antigen during vaccine production.

Potency is key parameter for evaluating the vaccine that limit the spread of disease. According to European Pharmacopeia (Monograph 04/2005:0063), vaccine potency is determined by measuring the protective dose through challenge studies in animals vaccinated with different antigen payloads. Protective dose 50 (PD₅₀) value of vaccine is well-defined by number of protective doses of antigen which protects fifty percent vaccinated population, when challenged with live virus. Value of PD₅₀ equal or greater than 3 is considered as acceptable for routine vaccination, while equal or greater than 6 is recommended in emergency vaccination in the case of FMD [17,18].

Despite being the gold standard, in-vivo potency test is highly expensive experiment, labor intensive, and associated with ethical concern in FMD endemic countries like Pakistan. These experiments require maintaining high bio-containment animal facilities to safeguard the premises from challenging infection, availability of susceptible host and involve exposure of animals to virulent virus. Furthermore, approximately 50% of unprotected animals may develop painful clinical manifestation of the disease. Even the animals with protective immunity will show lesions at live virus inoculation site and reduced in appetite and weight [19].

Consequently, there is growing interest in adopting substitute techniques that lessen the usage of animals and uphold the 3Rs: replacement, reduction, and refinement [20]. As alternatives to challenge-based potency testing, serological tests like the virus neutralization test (VNT) and enzyme-linked immunosorbent assay (ELISA) have been thoroughly studied. ELISA and VNT significantly lessen animal suffering while estimating the expected level of protection by evaluating antibody responses against virulent pathogen. Potency of FMD vaccine could be estimated indirectly by serological tests capable

of detecting neutralizing antibodies against structural proteins of FMDV [21]. Neutralizing antibodies directed against the structural proteins of FMDV, particularly intact 146S particles, are recognized as an important correlate of protection and can provide valuable information regarding vaccine performance. However, FMD protection is influenced by multiple factors, including antigenic matching between vaccine and field strains, vaccine formulation, adjuvant composition, duration of immunity, and cell-mediated immune responses. Therefore, although serological assays are valuable tools for monitoring immune responses and predicting vaccine efficacy [22], they cannot fully replace challenge-based potency studies, which remain essential for definitive evaluation of FMD vaccine protection

So therefore, to gain access to international market for the export of FMD free livestock, meat and other commodities, it is important to ensure that vaccines used in FMD control program/campaign should meet the purity requirement with regards to their NSP contents and it contain at least minimum required Protective Doses fifty (3PD₅₀). The study was design to formulate monovalent purified vaccine of local serotype A, Asia1 and O of FMDV with different concentration to estimate PD₅₀ as well as to evaluate anti NSP antibodies.

MATERIAL AND METHODS

Ethical Approval

All animal experiments were reviewed and approved by the Ethical Review Committee, Office of Research Innovation and Commercialization, University of Veterinary and Animal Sciences (approval number: No.DR/157, April 29, 2025), and conducted in accordance with the institutional guidelines for the care and use of laboratory animals.

Sample Collection and Confirmation

Vaccine serotype 'A, Asia 1, O' of FMDV was collected from Quality Operation laboratory (QOL), University of Veterinary and Animals Sciences (UVAS) Lahore. The viruses were confirmed through antigen detection ELISA kit (FMDV antigen and serotyping ELISA, the Pirbright Institute UK, Lot# 01-2021210708a).

Revival of the Cells

Cryopreserved baby hamster kidney (BHK-21) cell line was obtained from Quality Operation Laboratory, University of Veterinary and Animal Sciences. The cells were revived by propagating in Glasgow's Minimal essential medium (GMEM) (Cassion USA) with 10% fetal bovine serum (FBS) at 37°C for 24 h incubation in T-25 flask with 5% CO₂. Next day the cells were disintegrated through trypsin (Gibco, 0.25%) to check the viable count

by dye exclusion method (Trypan blue) and 10⁶ cells/ml were seeded into T-275 flasks and again the incubated at the above condition to develop monolayer. After 24 h cells were checked under inverted microscope for 80-90% monolayer.

Virus Propagation

After the establishment of confluent monolayer (80-90%) in flasks the media was removed. The cells were washed with sterile Phosphate Buffer Saline at least three times. The labeled cells culture flasks were inoculated with filtered FMDV serotype O, A and Asia separately under aseptic conditions for 37°C/40 min to adsorb the virus on cell surface. The flasks were seeded with infection medium (GMEM + 1% FBS) and incubated at 37°C for 24 h (O serotype), 36 h (A & Asia serotype). Following incubation time Cytopathic effects (CPEs) were observed under inverted microscope. After optimal CPEs the flasks were freeze thaw 2-3 time for the release of FMDV from cells. The suspensions containing the FMDV were centrifuge at 1000 rpm/10 min and supernatant were collected and stored at -40°C for further used.

Biological Titration

The stored FMDV serotypes were serially diluted tenfold separately in medium with 1% FBS. Three 96 well plates with monolayer of cells were prepared. The exhausted medium was discarded and each plate was inoculated 100 µL of dilated FMDV in respective wells and up to 10th well while 11 well with pure virus and 12 well with cell control. The plates were incubated at 37°C in CO₂ incubator. Following the incubation CPEs were recorded in 50% as an end point. The biological titer of FMDV serotypes were recorded as a 10_{log} Tissue culture infective dose in samples of FMDV as described by; 50% endpoint titer (TCID₅₀/0.1 mL) = 10_{log} total dilution above 50% + [PD*log (dilution factor)]

Inactivation of Virus

Clarified FMDV were chemically inactivated by using final concentration 3 mM binary ethyleneimine (BEI, MP Biomedical) and 0.04% formaldehyde (MP Biomedical). The residual of inactivation agents were neutralized by 2% Sodium Bis sulphate and sodium thiosulphate for BEI and Formaldehyde respectively. Sample from each inactivated virus were subjected to BHK-21 cells culture flask to check the live viral particle and incubated at 37°C for 24 & 36 h.

Concentration of Antigen Through Polyethylene Glycol (PEG-6000)

Inactivated FMDV serotype A, Asia 1 and O was divided two parts. One part of each serotype was concentrated using PEG-6000 (Deajung, Korea). 100 mL of each virus was inoculated 7.5% PEG from 50% concentrated

solution (Tris Buffer, pH 7.2) and suspension was placed at stirring at 4°C. The suspension was centrifuged and pelleted. Supernatant was discarded and elution of antigen was done using PBS (0.011M, pH 7.2). 1/10 of the original volume of antigen of each virus was obtained and stored at 4°C for further action.

Purification and Quantification of FMDV by Size Exclusion Chromatography (SEC)

The antigen was purified and quantified using BioRad Biologic Chromatograph system 358-BR3506 with UV monitor (2 mm path length) at 254 nm. Biorad Econo Column model 15/50 was filled with resin Sephacryl™ S-300 high resolution (Cytiva, Germany) using PBS 0.011M pH 7.2 and resolution was monitored using solution of Blue dextran (BD) and Bovine Serum Albumen (BSA) with optimized parameter like flow rate (FR = 0.75 mL/min), bed height (48cm), sample volume (W=3.5 mL), path length of flow cell (PL = 0.2 cm), chart recorder speed 12 cm/h (S = 0.2 cm/min). The concentrated sample of each virus was run on chromatographic system and eluted was collected and stored at 4°C for vaccine preparation used. The chromatogram (peaks) of each virus was recorded 254 nm with $E_{cm1\%} = 72$; different sensitivity absorbance unit (FSD) used for the estimation of FMDV in µg/mL using formulae;

$$\text{FMDV in sample } (\mu\text{g/mL}) = \frac{\text{FR} \times \text{PA} \times \text{FSD} \times 1000}{\text{S} \times \text{PL} \times \text{E} \times \text{W}} \quad [23]$$

Purity Confirmation of FMDV

The chromatographic elute of FMDV was run on Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). For this purpose, resolving gel and stacking gel was prepared at concentration 15% & 5% respectively. Sample of chromatographic elute and concentrated was loaded with prestained protein marker (10-180kDa, Servicebio, China) onto well and with Tris-Glycine buffer at 190V for continuous current. The size of band pure and crude FMDV was compared with protein marker.

Vaccines Formulation and Quality Control

Monovalent purified inactivated vaccine of each serotype with different dilution was prepared. For this purpose, 50% of aqueous phase containing quantified contents of 146S fraction of FMDV with 50% of montanide oil ISA 50 (France) as adjuvant was emulsified for each antigen. Five different dilutions of each virus (A' serotype as A₁ A₂ A₃ A₄ A₅), (Asia 1' as B₁ B₂ B₃ B₄ B₅) & ('O' as C₁ C₂ C₃ C₄ C₅) at 16, 8, 4, 2 & 1 µg/dose was adjusted with final dose volume 2 mL sterile PBS. Thiomersal sodium was added at 0.01% as preservative. Each vaccine was mixed thoroughly with sterilized homogenizer at room temperature to get consistent vaccine suspension. The prepared purified inactivated vaccine was evaluated for its safety, sterility,

Innocuity and potency testing as described by Razak et al.^[9]. Physical parameters of the vaccines like viscosity, stability and emulsion type were validated as described by Shabana and Abd EL-Sadek^[24].

Selection of Calves and Inoculation of Vaccines and Serum Collection

A total of seventy-five healthy, adult and unvaccinated bovine calves (above 5 months of age, 25/group, 5/ dilution/vaccine type) of randomly selected breed raised at private cattle farm at FMD free region Bahawalpur, Punjab Pakistan were selected randomly and used for the estimation of potency testing of purified monovalent FMD vaccines. The calves were screened for the presences of antibodies against FMDV type through solid phase competitive ELISA (SPCE). The animals found non protective against FMDV were selected for potency testing and divided into four main group A, B, C & D with each subgroup for each dilution ('A' serotype as A₁ A₂ A₃ A₄ A₅), ('Asia 1' as B₁ B₂ B₃ B₄ B₅) & ('O' as C₁ C₂ C₃ C₄ C₅) and D (Control group, 2/each group). Each group was inoculated with primary dose and following 28 days with booster dose of vaccine from each dilution. All animals of each group were kept under observation for five days post vaccination. Any kind of allergic reaction, edema formation, erythema, changes in body temperature, change in feed intake or any other symptoms were observed. The serum of all animals was collected after 28 days of booster inoculation and tested for the presence of antibodies.

Solid Phase Competitive ELISA (SPCE)

The commercially available blocking ELISA (PrioCHECK™ FMDV Type A, Type Asia 1 and Type O Ab Strip Kit) was used for the detection of antibodies against FMDV. Each ELISA micro plate was coated with non-infectious FMDV type antigen. The reaction between FMDV type and a specific monoclonal antibody (mAb) was blocked by specific antibodies that were present in the test serum of bovine calves and rests of the steps were performed according instruction. For serotype (A, Asia 1), $PI = 100 - (\text{OD}_{450} \text{ sample} / \text{OD}_{450} \text{ Max.}) * 100$ ($\text{OD}_{450} \text{ max} = \text{control serum (Mean OD} \geq 1.00)$). For serotype O, $PI = 100 - (\text{Corrected OD}_{450} \text{ sample} / \text{Corrected OD}_{450} \text{ Max.}) * 100$ (Whereas $\text{Corrected OD}_{450} \text{ sample} = (\text{OD}_{450} \text{ sample} - \text{Blank}) / \text{Corrected OD}_{450} \text{ Max}$ is control serum) $PI < 50\%$ is considered negative for FMDV while $PI \geq 50\%$ is considered positive for FMDV.

Determination of Mass Load of FMDV and Estimation of Protective Dose 50 (PD₅₀)

The test sera of animals vaccinated with FMD vaccine having five different concentrations of the 146S fractions for each serotype were declared protective or non-protective based upon the percent inhibition of solid

phase competition ELISA. The minimum concentration of the 146S fraction which elicits 100% protective immune response in calves was determined. Apart from this number of protective doses 50 (PD₅₀) of monovalent FMDV vaccine containing minimum concentration of FMDV Ag but eliciting 100% protection in group of animals based upon SPCE according to WOAHA was estimated as described by Spearman-Kärber's method.

$$PD_{50} (\log_{10}) = - [X_0 - D/2 + D (\sum R/N)]$$

Where, X₀ is log₁₀ of reciprocal of lowest concentration of antigen at 100% protection; D is the log 10 of dilution factor; N is total animals in each group; R is protected out of total animals.

Trivalent Vaccine Formulation and Protective Immune Response

Trivalent vaccine (3PD₅₀) containing purified antigen Serotype A, Asia 1 & O with concentration 6.42 µg, 5.7 µg and 5 µg respectively was formulated with 50% montanide ISA 50 oil adjuvant with standard protocol by adjusting final volume 2 mL/dose. After quality control testing, vaccine was inoculated to five healthy, non-vaccinated calves with primary and booster dose at day 0, and 4th week. Serum was collected at 4th week post booster dose and evaluated for protective antibody titer through Solid Phase competitive ELISA. Anti-NSP antibodies were measured through commercial ELISA kit (IDEXX, 3ABC Bov-Ov). NSP < 20% & ≥ 30% were subjected/considered negative and positive respectively.

Statistical Analysis

The percentage inhibition (PI) values obtained from SPCE ELISA for serotypes A, Asia 1, and O at day 0, 4th week, and 8th week post-vaccination were statistically analyzed by repeated measure ANOVA using SPSS (Version 20, IBM Corp., USA).

RESULTS

Confirmation of Serotypes

FMDV serotyping was confirmed by antigen detection ELISA with positive (≥1.0) and negative control (≤0.1). All the sample were serotyped in their respective well and value for consider to be positive must be greater than 0.1. And for the test validity positive well must be ≥1.0 while negative must be ≤0.1 (Table 1).

Cells Monolayer Development and Propagation of Virus

Cells were revived from stock under aseptic condition and monolayer of cells was first developed in T-25 flask. Following the process the viable cells were counted through dye exclusion method and cells were transferred to T-275 flasks for monolayer developed. After specific incubation

time the 80-90% developed monolayers on specified flask were selected as shown in Fig. 1-A, C, E. Serotype A, Asia 1 & O of FMDV was propagated and incubated at 37°C for 18-24 h & 36 h for O and A, Asia serotype respectively for replication of virus. As virus replicate the specific changes on cells were observed in term of cytopathic effects as shown in Fig. 1-B, D, F). After completion of incubation and observation of CPEs the flasks were freeze thaw for 2-3 time and centrifuge to collect supernatants and stored at -20°C.

Tissue Culture Infective Dose 50 Calculation of Serotype A, Asia 1 and O

Biological titration in term of Tissue Culture Infective Dose 50 (TCID₅₀) for quantification of FMDV serotype A, Asia & O to produce fifty percent cytopathic effects on BHK-21 cells were estimated through reed and murch method (Table 2).

Assessment of Inactivation and Concentration

Inactivated sample of each serotype was inoculated on BHK-21 cells to evaluate the inactivation of the viral antigen. Repeatedly seven blind passages were given to each serotype and CPEs were checked. Up to the 7th

Table 1. serotyping of FMDV through Antigen Detection ELISA

Catching MABs	Sample	POS Control	NEG Control
Type O	0.545	1.531	0.0332
Type A (1 st MAB)	0.940	1.531	0.0332
Type A (2 nd MAB)	0.031	1.531	0.0332
Type Asia 1	0.297	1.531	0.0332
Type C	0.027	1.531	0.0332
Pan O, A, C & Asia 1	0.517	1.531	0.0332




Fig 1. (A) confluent monolayer of BHK-21 Cells, (B) CPEs caused by FMDV serotype 'A' infection (C) confluent monolayer of BHK-21, (D) CPEs caused by FMDV serotype 'Asia 1' infection (E) confluent monolayer of BHK-21, (F) CPEs caused by FMDV serotype 'O' infection

Table 2. Calculation of biological titration (TCID₅₀)

Serotype	Dilution	Wells with CPE	Wells without CPE	CP	CN	Total	% CPE	TCID ₅₀ /mL
Serotype A	10 ⁻¹	8	0	46	0	46	100	10 ^{7.29}
	10 ⁻²	8	0	38	0	38	100	
	10 ⁻³	8	0	30	0	30	100	
	10 ⁻⁴	7	1	22	1	23	95.65	
	10 ⁻⁵	6	2	15	3	18	83.33	
	10 ⁻⁶	5	3	9	6	15	60.00	
	10 ⁻⁷	3	5	4	11	15	26.66	
	10 ⁻⁸	1	7	1	18	19	5.26	
	10 ⁻⁹	0	8	0	26	26	0.00	
	10 ⁻¹⁰	0	8	0	34	34	0.00	
Serotype Asia 1	10 ⁻¹	8	0	49	0	49	100	10 ^{7.61}
	10 ⁻²	8	0	41	0	41	100	
	10 ⁻³	8	0	33	0	33	100	
	10 ⁻⁴	8	0	25	0	25	100	
	10 ⁻⁵	7	1	17	1	18	94.44	
	10 ⁻⁶	6	2	10	3	13	76.92	
	10 ⁻⁷	3	5	4	8	12	33.33	
	10 ⁻⁸	1	7	1	15	16	6.25	
	10 ⁻⁹	0	8	0	23	23	0.00	
	10 ⁻¹⁰	0	8	0	31	31	0.00	
Serotype O	10 ⁻¹	8	0	45	0	45	100	10 ^{7.19}
	10 ⁻²	8	0	37	0	37	100	
	10 ⁻³	8	0	29	0	29	100	
	10 ⁻⁴	7	1	21	1	22	95.45	
	10 ⁻⁵	6	2	14	3	17	82.35	
	10 ⁻⁶	5	3	8	6	14	57.14	
	10 ⁻⁷	2	6	3	12	15	20.00	
	10 ⁻⁸	1	7	1	19	20	5.00	
	10 ⁻⁹	0	8	0	23	23	0.00	
	10 ⁻¹⁰	0	8	0	31	31	0.00	

passage no development of CPEs indicated that viruses were inactivated successfully. Following the procedure the 100 mL of each serotype were concentrated 10X (1/10th of original) through PEG-6000 to obtain enough antigens after purification through SEC as it dilutes the sample. About 10ml of each concentrated virus was run on chromatography.

Purification and Quantification Through SEC and SDS-PAGE Analysis

Each concentrated sample was run on chromatographic system and elute was collected. Chromatogram of each virus (Fig. 2-A, B, C) was obtained and peak area was calculated to quantify the virus in sample and eluted in

term of µg/mL in as described in Table 3. Elute of the sample A and Asia 1 was run on SDS-PAGE to check the purity. Structural proteins 64kDa (VP2=VP4), 25kDa (VP3) and 35kDa (VP1) and no bands of non-structural protein at 53kDa (3ABC) and 56kDa (3D) with respect to prestained protein marker (10-180x, Servicebio, China) for purifies FMDV. While in crude sample NSP bands at 53-56kDa along with structural protein were observed (Fig. 3). It is resulted that purified FMDV antigen were free from NSP contents.

Preparation of Monovalent Vaccines

The purified and quantified elute of antigens were diluted to prepared stock of 16 µg/mL of each virus




Table 3. Chromatographic attributes of FMDV serotypes and antigen load ($\mu\text{g}/\text{mL}$)			
Chromatographic Attributes	Serotype A	Serotype Asia 1	Serotype O
Flow rate of mobile phase (FR)	0.7 mL/min	0.7 mL/min	0.7 mL/min
Sample volume loop (W)	3.5 mL (4% of CV)	3.5 mL (4% of CV)	3.5 mL (4% of CV)
Speed of chart recorder (S)	12 cm/h	12 cm/h	12 cm/h
Extinction point of FMDV ϵ	72	72	72
Path length of flow cell (PL)	0.2 cm	0.2 cm	0.2 cm
Full scale absorbance unit setting (FSD)	0.3	0.5	0.3
Elution volume	28 mL (40 x 0.7)	37.5 mL (54 x 0.7)	37.5 mL (54 x 0.7)
Dilution factor (DF)	28/3.5 = 8	37.5/3.5 = 10.8	37.5/3.5 = 10.8
Length of peak	5.4 cm	5.4 cm	5.0 cm
Base of peak	4.5	5	4.7
Area under peak	12.37 cm^2	13.5 cm^2	11.75 cm^2
Crude concentration * =	= 257	= 468	= 244
Elute Concentration ($\mu\text{g}/\text{mL}$) = Crude/Dilution factor	257/8 = 32.125	468/10.8 = 43.75	244/10.8 = 22.59

(Table 4). The purified monovalent vaccines (five concentration of each antigen) were prepared by emulsifying montanide oil ISA 50 and adjusted volume 2 mL/dose (15 doses/30 mL) with sterile PBS as shown in table. Quality testing was performed as per standard protocol.

Vaccine Inoculation and Animal Response

Vaccine of each antigen/concentration was inoculated

to five bovine calves at day 0 & 21st (booster) and blood was collected at different interval day 0, 4th and 8th week of vaccination. Serum was collected and checked through Solid Phase Competitive ELISA for calculation of percentage inhibition (PI) (PI \geq 50% protected, PI<50% non-protected) as shown in Table 5. A P-value of less than 0.05 in Mauchly's test of sphericity indicated that the assumption of sphericity was violated, demonstrating significant differences in




Fig 3. 64kDa (VP2+VP4), 25kDa (VP3) and 35kDa (VP1) Bands of purified FMDV serotypes on SDS-PAGE after elution from Size Exclusion Chromatography (SEC)

Table 4. Formulation of Purified Inactivated Monovalent FMD Vaccines, against A, Asia 1 and O serotype (Monovalent vaccine for each serotype with same layout)

B #	146S Fraction (µg)/Dose	Ingredients (mL)				Total Volume of Vial/Dose
		Antigen	Montanide Oil	Thiomersal Sodium (µL)	PBS	
B1	16	15	15	3	--	30/15
B2	8	7.5	15	3	7.5	30/15
B3	4	3.75	15	3	11.25	30/15
B4	2	1.875	15	3	13.125	30/15
B5	1	0.9375	15	3	14.065	30/15

percentage inhibition (PI) across day 0, week 4, and week 8. Furthermore, the time × serotype interaction effect was also statistically significant ($P < 0.05$), indicating that the pattern of antibody response over time differed significantly among serotypes A, Asia 1, and O.

Table 5. PI of SPCE for Serotype A, Asia 1 and O, at day zero, 4th & 8th week

Group A (Serotype A)				Group B (Serotype Asia 1)				Group C (Serotype O)			
Animal Tag No.	Day 0	4 th Week	8 th Week	Animal Tag No.	Day 0	4 th Week	8 th Week	Animal Tag No.	Day 0	4 th week	8 th Week
A ₁₁	41.53	58.78	75.31	B ₁₁	26.52	56.14	82.84	C ₁₁	23.97	55.72	84.57
A ₁₂	31.62	50.95	69.88	B ₁₂	33.15	54.55	79.39	C ₁₂	26.20	51.73	81.41
A ₁₃	40.97	69.72	72.76	B ₁₃	38.19	52.69	88.94	C ₁₃	20.04	53.18	83.64
A ₁₄	44.32	57.58	76.91	B ₁₄	39.25	61.27	87.53	C ₁₄	22.11	52.56	78.51
A ₁₅	41.29	61.42	78.19	B ₁₅	49.04	57.02	88.94	C ₁₅	27.34	50.28	85.60
A ₂₁	36.26	78.99	74.52	B ₂₁	38.28	55.87	83.28	C ₂₁	19.78	53.96	77.68
A ₂₂	30.99	67.41	72.44	B ₂₂	40.40	59.23	90.80	C ₂₂	15.27	53.75	81.67
A ₂₃	33.62	56.94	62.38	B ₂₃	43.85	57.73	87.97	C ₂₃	23.14	53.08	76.44
A ₂₄	36.82	66.21	74.92	B ₂₄	38.01	59.85	90.18	C ₂₄	21.23	52.56	82.08
A ₂₅	38.89	61.74	76.11	B ₂₅	32.36	65.07	82.40	C ₂₅	32.72	50.70	82.81
A ₃₁	35.06	58.14	67.49	B ₃₁	34.30	61.62	81.16	C ₃₁	25.16	53.65	82.96
A ₃₂	21.00	51.27	53.43	B ₃₂	39.08	54.46	85.05	C ₃₂	22.26	37.86	45.83
A ₃₃	32.74	60.14	78.43	B ₃₃	28.20	62.68	88.15	C ₃₃	27.60	50.08	73.54
A ₃₄	38.97	48.80	49.28	B ₃₄	31.83	57.73	86.03	C ₃₄	18.12	50.28	77.01
A ₃₅	28.03	62.85	82.82	B ₃₅	38.90	45.88	48.98	C ₃₅	27.03	54.89	84.41
A ₄₁	14.61	47.36	48.16	B ₄₁	34.92	59.77	84.52	C ₄₁	17.55	56.60	36.30
A ₄₂	25.31	58.22	85.14	B ₄₂	26.17	37.04	38.19	C ₄₂	29.67	51.42	74.31
A ₄₃	47.36	42.65	46.24	B ₄₃	41.73	58.53	86.29	C ₄₃	15.06	52.46	76.44
A ₄₄	8.86	44.16	46.96	B ₄₄	39.52	63.92	85.05	C ₄₄	22.06	27.91	48.99
A ₄₅	41.45	55.43	73.16	B ₄₅	35.19	47.03	48.18	C ₄₅	9.94	51.89	72.14
A ₅₁	32.26	57.90	65.73	B ₅₁	26.79	33.24	36.51	C ₅₁	29.41	37.86	41.22
A ₅₂	23.64	46.40	49.28	B ₅₂	41.99	46.41	46.15	C ₅₂	14.39	53.34	70.17
A ₅₃	34.10	30.35	46.00	B ₅₃	36.25	40.84	40.40	C ₅₃	22.09	51.73	59.14
A ₅₄	25.07	48.00	49.12	B ₅₄	34.74	55.08	68.70	C ₅₄	19.26	23.25	35.53
A ₅₅	28.19	44.48	47.84	B ₅₅	38.19	45.62	47.39	C ₅₅	12.42	27.29	33.66
Control	37.69	38.55	38.01	Control	34.10	35.54	36.02	Control	29.47	35.68	30.50
	38.54	40.31	44.82		38.33	39.88	43.92		33.45	38.58	37.55

Table 6. Estimation of protective dose (PD₅₀)

Serotype	G. ID	146S con. (µg/dose)	Dilution	PC	NPC	CPC	CNPC	Total Animals	Protection %	Min. Con. of Ag. (µg/dose) for 50% Protection
Serotype A	A ₁	16	Stock	5	0	17	0	17	100	2.14
	A ₂	8	1:2	5	0	12	0	12	100	
	A ₃	4	1:4	4	1	7	1	8	87.5	
	A ₄	2	1:8	2	3	3	4	7	42.85	
	A ₅	1	1:16	1	4	1	7	8	12.5	
$PD_{50}(\text{Log}10) = [0 - 0.3010/2 + 0.3010 (12/5)] = 0.5719 = 3.73 PD_{50}/\text{dose}$										
Serotype Asia 1	B ₁	16	Stock	5	0	18	0	18	100	1.90
	B ₂	8	1:2	5	0	13	0	13	100	
	B ₃	4	1:4	4	1	8	1	9	88.8	
	B ₄	2	1:8	3	2	4	3	7	57.14	
	B ₅	1	1:16	1	4	1	7	8	14.28	
$PD_{50}(\text{Log}10) = [0 - 0.3010/2 + 0.3010 (13/5)] = 0.6321 = 4.21 PD_{50}/\text{dose}$										
Serotype O	C ₁	16	Stock	5	0	19	0	19	100	1.63
	C ₂	8	1:2	5	0	14	0	14	100	
	C ₃	4	1:4	4	1	9	1	10	90.0	
	C ₄	2	1:8	3	2	5	3	8	62.50	
	C ₅	1	1:16	2	3	2	6	8	25.5	
$PD_{50}(\text{Log}10) = [0 - 0.3010/2 + 0.3010 (14/5)] = 0.6923 = 4.92 PD_{50}/\text{dose}$										

PC: protected calves, NPC: non-protected calves, CPC: cumulative protected calves, CNPC: cumulative non-protective calves

Estimation of PD₅₀

On the basis of ELISA percentage inhibition interpretation, the animals declared protective (P) and non-protective (NP). Protective Dose 50 for FMDV type A was measure as lowest concentration of antigen (8 µg/dose) give 100 percent protection (PD₅₀≥3). Likewise, PD₅₀ for Asia 1 and O was determined & describe in *Table 6*, respectively.

Trivalent Vaccine Preparation, Inoculation and Immune Response

Trivalent vaccine (equal to 3PD₅₀) was prepared by containing antigen concentration (6.42, 5.7 and 4.90 µg/dose) as calculated by estimation of protective dose 50. The five animals were inoculated at day 0 and 4th week (booster dose). Blood was collected at different interval and serum was obtained to check the combined effect of trivalent vaccine and PI was calculated through SPCE. All Animals were protected against three antigens (*Table 7*). Anti NSP antibodies were checked to assess the purified potential of the vaccine through NSP ELISA (IDEXX, 3ABC Bov-Ov). The entire animal population showed <20% NSP antibodies contend (*Table 7*).

Table 7. Trivalent vaccine (3PD₅₀) evaluation through SP & NSP ELISA

Time Interval	Percentage Inhibition for SPCE			
	A	Asia1	O	Control
Day zero	38.18±10.16	37.79±8.16	30.16±6.61	27.11±6.51
4 th week	62.49±4.90	65.96±4.29	57.71±2.25	29.44±6.23
8 th week	79.28±6.95	83.11±7.43	84.97±4.77	32.99±4.42
Percentage for 3ABC NSP				
Day zero	8.83±1.16			
4 th week	8.01±0.77			
8 th week	7.78±1.67			

DISCUSSION

FMD is devastating disease of animals which produced heavy economic losses in term of meat, milk and related products. These products have restriction on export to disease free countries. Potent vaccine is ultimate need to control the disease and spread of live virus. The endemic countries need to check the potency of vaccine to produce NSP free vaccine with DIVA strategy (to distinguish vaccinated from infected animals) to take access to global market for their animal products. To evaluate the potency of the vaccine in-vivo by challenging animals is an

expensive task. Scientists are agreed upon Replacement, Reduction and Refinement (3R) concept to reduce the challenge phase of the testing for animal welfare and cost issues. Alternative methods like challenge free serological approaches are being used with relation of antibody titer to protection level [25].

ELISA OD values for A, Asia1 & O were 0.545, 0.940, and 0.297 respectively (≥ 0.1) [26]. The cells monolayer with 80-90% was developed on flat bottom flasks with inoculation of 10^6 cells/mL at 37°C for 24-36 h. The virus was propagated and after CPEs development virus was harvested and biological titer $10^{7.29}$, $10^{7.61}$, $10^{7.19}$ TCID₅₀ for serotype A, Asia1 & O was calculated using reed and much method. The results were consistent accordance with the study by study [27]. Harvested culture of each serotype virus was double inactivated by BEI and formaldehyde with 3mM and 0.04% respectively followed by ensuring the inactivation process by inoculation on cell culture upto the 7 blind passages. The results are supported by investigation [28].

SEC is efficient technique to purify the 146S content of the FMDV from nonstructural and media protein. Sephacryle S-300 have exclusion limit 10kDa-1500kDa and all the protein from media and NSP are trapped while FMDV (8200kDa) can pass between resin beads; therefore, eluted rapidly [16]. Peak area from UV detector in the form of chromatogram is obtained that is directly related to virus concentration. The optimized condition according to the referenced article [16] were used to calculate the virus concentration 32.125, 43.75 and 22.59 $\mu\text{g/mL}$ for A, Asia1 and O serotype antigen respectively. The result are supported by the study by [16]. Elute was run on SDS-PAGE to check the purity of FMDV with crude virus as a control. Bands of 64, 35, & lighter 25kDa of VP2+VP4, VP1 and VP3 respectively of purified FMDV were seen using 10-180x prestained protein marker. While in crude FMDV the 53-56kDa band of 3ABC & 3D was seen which was absent in the Purified product. These findings are accordance to study which use purified FMDV in SDS-PAGE from SDG method [29].

The vaccine containing 16 & 8 $\mu\text{g/dose}$ of A serotype showed 100% protection while 4, 2 and 1 $\mu\text{g/dose}$ animal group were 87.5, 42.85, & 12.5% respectively. The protective dose in group of animals with 100% percent protection and lowest concentration of virus (8 $\mu\text{g/dose}$) while was 3.73PD₅₀/dose. These are supported by study [9]. Minimum concentration of 2.14 μg was calculated which protect 50% animal in the group. For A serotype 3PD₅₀ = 6.42 $\mu\text{g/dose}$ was estimated in this study. 100% animal protection was observed in vaccine type Asia1 with concentration 16 & 8 $\mu\text{g/dose}$ while remaining 4, 2 & 1 $\mu\text{g/dose}$ were 88.8,

57.14, and 14.28% that mean lowest concentration of 8 $\mu\text{g/dose}$ showed 100% percent protection with PD₅₀ = 4.21/dose. Minimum concentration for 1PD₅₀ was 1.90 $\mu\text{g/dose}$ while 3PD₅₀ = 5.7 $\mu\text{g/dose}$. In the case of O serotype 100% animal protection were calculated in 16 & 8 $\mu\text{g/dose}$; however, the 4, 2, and 1 $\mu\text{g/dose}$ showed the 90.0, 62.5 & 25.5% protection with calculated as 4.92PD₅₀/doses of the lowest concentration of 8 $\mu\text{g/dose}$ (100% protection). Estimated minimum concentration conferring PD₅₀ was 1.63 $\mu\text{g/dose}$ with 3PD₅₀ = 4.89 $\mu\text{g/dose}$. These above findings are accordance with studies [19,30,31] $\geq 3\text{PD}_{50}$ is international requirement for routine vaccine and 6PD₅₀ for emergency vaccination [32].

Trivalent vaccine of FMDV equal to 3PD₅₀ concentration 6.42, 5.7, and 4.90 $\mu\text{g/dose}$ was prepared and inoculated to healthy, with no prior vaccine history of animals (n=5) with primary and booster dose at day zero and 28th week. After 4th week of booster dose serum was collected and checked for SPCE & NSP ELISA. All the animals were protective at week 4 & 8 to three serotypes with mean PI for serotype A; 62.49 \pm 4.90 & 79.28 \pm 6.95, Asia1 65.96 \pm 4.29, 83.11 \pm 7.43 and O 57.71 \pm 2.25, 84.97 \pm 4.7 while NSP contents in all animals were (Mean \pm SD) 8.83 \pm 1.16, 8.87 \pm 0.77, 7.78 \pm 1.67 at day zero, week 4&8 respectively, were less than the cutoff value (20%). Same NSP-ELISA Approach was used and results are accordance with referenced study [33].

In conclusion, present study determined the PD₅₀ values of chromatographically purified FMDV serotypes A, Asia1, and O using serological techniques and evaluated their application in the formulation of a trivalent vaccine in bovines. These findings demonstrate that serological techniques can be alternatively applied for the estimation of PD₅₀ values of purified FMD vaccines, providing a practical approach for vaccine potency assessment and antigen payload standardization in resource setting areas. Furthermore, the study supports the use of SEC as an efficient method for the production of purified FMD vaccines with reduced NSP content and improved consistency for vaccine formulation.

DECLARATIONS

Availability of Data and Materials: The data supporting the findings of this study are available from the corresponding author (I.A.) upon reasonable request.

Acknowledgement: The authors are grateful to all members of the Institute of Microbiology, UVAS, Lahore, Pakistan, for their support in conducting this work. Special acknowledgment is extended to the late Prof. Dr. Aftab Ahmad Anjum for his valuable contributions to the study design and guidance. His scientific insight and mentorship are sincerely appreciated and remembered.

Financial Support: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical Approval: All animal experiments were reviewed and approved by the Ethical Review Committee, Office of Research Innovation and Commercialization, University of Veterinary and Animal Sciences (approval number: No.DR/157, April 29, 2025).

Conflict of Interest: The authors declare that they have no relevant financial or non-financial interests to disclose.

Declaration of Generative Artificial Intelligence (AI): The authors declare that no artificial intelligence or AI-assisted technologies were used in the preparation of this article, including the creation of text, tables, or figures except to improve the readability and language of the article.

Author Contributions: FA and IA conceptualize, designed, executed the experiment, compiled the results and wrote the manuscript. ARA analyzed the results. AG interpreted the results, critically revised the manuscript and approved for submission.

REFERENCES

- Zahur A, Irshad H, Hussain M, Anjum R, Khan M: Transboundary animal diseases in Pakistan. *J Vet Med B*, 53 (Suppl. 1): 19-22, 2006. DOI: 10.1111/j.1439-0450.2006.01015.x
- Harmsen MM, Li H, Sun S, Van Der Poel WH, Dekker A: Mapping of foot-and-mouth disease virus antigenic sites recognized by single-domain antibodies reveals different 146S particle specific sites and particle flexibility. *Front Vet Sci*, 9:1040802, 2023. DOI: 10.3389/fvets.2022.1040802
- Dhakal SP, Pandey K, Upadhyaya M, Karki S, Ramdam N, VanDyk S, Premashthira S: Spatiotemporal distribution of foot-and-mouth disease in Nepal between 2019 and 2021. *Anim Dis*, 3 (1):39, 2023. DOI: 10.1186/s44149-023-00104-9
- Islam M, Akter M, Hassan M, Rahman M, Islam E, Khan M, Chakrabarty A, Giasuddin M: Identification of Foot and Mouth Disease (FMD) virus from recently outbreak crossbred cattle in Rajbari District of Bangladesh. *SAARC J Agric*, 19 (1): 201-210, 2021. DOI: 10.3329/sja.v19i1.54790
- Al-Husseiny SH, Kshash QH, Jassim A: Sero-detection of foot and mouth disease virus serotypes A and O in one-humped camels (*Camelus dromedarius*) in the Middle of Iraq. *Kafkas Univ Vet Fak Derg*, 26 (6): 743-747, 2020. DOI: 10.9775/kvfd.2020.24276
- Stenfeldt C, Eschbaumer M, Pacheco JM, Rekant SI, Rodriguez LL, Arzt J: Pathogenesis of primary foot-and-mouth disease virus infection in the nasopharynx of vaccinated and non-vaccinated cattle. *PLoS One*, 10 (11):e0143666, 2015. DOI: 10.1371/journal.pone.0143666
- Elrashedy A, Nayel M, Salama A, Zaghawa A, El-Shabasy RM, Hasan ME: Foot-and-mouth disease: Genomic and proteomic structure, antigenic sites, serotype relationships, immune evasion, recent vaccine development strategies, and future perspectives. *Vet Res*, 56 (1):78, 2025. DOI: 10.1186/s13567-025-01485-0
- Gao Y, Sun SQ, Guo HC: Biological function of foot-and-mouth disease virus non-structural proteins and non-coding elements. *Virol J*, 13 (1):107, 2016. DOI: 10.1186/s12985-016-0561-z
- Razak A, Altaf I, Anjum AA, Awan AR: Preparation of purified vaccine from local isolate of foot and mouth disease virus and its immune response in bovine calves. *Saudi J Biol Sci*, 30 (7):103709, 2023. DOI: 10.1016/j.sjbs.2023.103709
- Terry G, Clark RP, Rweyemamu M: Variations in the buoyant density of foot-and-mouth disease virus strains. *Arch Virol*, 71 (4): 333-341, 1982. DOI: 10.1016/j.sjbs.2023.103709
- Pons Royo MdC, Jungbauer A: Polyethylene glycol precipitation: Fundamentals and recent advances. *Prep Biochem Biotechnol*, 55 (8): 935-954, 2025. DOI: 10.1080/10826068.2025.2470220
- Kim AY, Park SY, Park SH, Kim JY, Jin JS, Kim E-S, Park JH, Ko YJ: Comparison of high-performance liquid chromatography with sucrose density gradient ultracentrifugation for the quantification of foot-and-mouth disease vaccine antigens. *Vaccines*, 10 (5):667, 2022. DOI: 10.3390/vaccines10050667
- Fernandez-Martinez J, LaCava J, Rout MP: Density gradient ultracentrifugation to isolate endogenous protein complexes after affinity capture. *Cold Spring Harb Protoc*, 2016 (7):pdb. prot087957, 2016. DOI: 10.1101/pdb.prot087957
- Goyon A, Beck A, Colas O, Sandra K, Guillarme D, Fekete S: Evaluation of size exclusion chromatography columns packed with sub-3 µm particles for the analysis of biopharmaceutical proteins. *J Chromatogr A*, 1498, 80-89, 2017. DOI: 10.1016/j.chroma.2016.11.056
- Hall M: Size exclusion chromatography (SEC). *Biopharmaceutical Processing: Development, Design, and Implementation of Manufacturing Processes*, 421-432, Elsevier, 2018.
- Razak A, Altaf I, Anjum AA, Awan AR, Awan FN: Optimization of size exclusion chromatography for the purification and quantification of foot and mouth disease virus serotype "O". *Pak J Zool*, 57 (1): 107-116, 2025. DOI: 10.17582/journal.pjz/20211109021143
- Waters R, Ludi AB, Fowler VL, Wilsden G, Browning C, Gubbins S, Statham B, Bin-Tarif A, Mioulet V, King DJ: Efficacy of a high-potency multivalent foot-and-mouth disease virus vaccine in cattle against heterologous challenge with a field virus from the emerging A/ASIA/G-VII lineage. *Vaccine*, 36 (14): 1901-1907, 2018. DOI: 10.1016/j.vaccine.2018.02.016
- Singh RP, Mohapatra JK: Review and evaluation of data defining FMD vaccine potency standards for endemic countries. *VirusDis*, 2026:1-10, 2026. DOI: 10.1007/s13337-026-00959-9
- Goris N, Maradei E, D'aloia R, Fondevila N, Mattion N, Perez A, Smitsaert E, Nauwynck H, La Torre J, Palma E: Foot-and-mouth disease vaccine potency testing in cattle using homologous and heterologous challenge strains: Precision of the "Protection against Podal Generalisation" test. *Vaccine*, 26 (27-28): 3432-3437, 2008. DOI: 10.1016/j.vaccine.2008.04.034
- Romberg J, Lang S, Balks E, Kamphuis E, Duchow K, Loos D, Rau H, Motitschke A, Jungbäck C: Potency testing of veterinary vaccines: The way from in vivo to in vitro. *Biologicals*, 40 (1): 100-106, 2012. DOI: 10.1016/j.biologicals.2011.10.004
- Paton D, Reeve R, Capozzo AV, Ludi A: Estimating the protection afforded by foot-and-mouth disease vaccines in the laboratory. *Vaccine*, 37 (37): 5515-5524, 2019. DOI: 10.1016/j.vaccine.2019.07.102
- Othuisit L, Kabelo TI, Nkwe KA, Seoke I, Fana EM, Lebani K: In vitro assessment of foot-and-mouth disease vaccine immunogenicity: Advances, challenges, and ethical considerations. *Front Vet Sci*, 13:1791118, 2026. DOI: 10.3389/fvets.2026.1791118
- Spitteler MA, Fernández I, Schabes E, Krimer A, Régulier EG, Guinzburg M, Smitsaert E, Levy MS: Foot and mouth disease (FMD) virus: Quantification of whole virus particles during the vaccine manufacturing process by size exclusion chromatography. *Vaccine*, 29 (41): 7182-7187, 2011. DOI: 10.1016/j.vaccine.2011.05.078
- Shabana WS, Abd EL-Sadek A: Effect of carbomer as an adjuvant for enhancement of immune-response against FMD vaccine. *Appl Vet Sci*, 6 (2): 27-36, 2021. DOI: 10.21608/jav.2021.154578
- Barnett PV, Statham RJ, Vosloo W, Haydon DT: Foot-and-mouth disease vaccine potency testing: Determination and statistical validation of a model using a serological approach. *Vaccine*, 21 (23): 3240-3248, 2003. DOI: 10.1016/s0264-410x(03)00219-6
- Grazioli S, Ferris NP, Dho G, Pezzoni G, Morris AS, Mioulet V, Brocchi E: Development and validation of a simplified serotyping ELISA based on monoclonal antibodies for the diagnosis of foot-and-mouth disease virus serotypes O, A, C and Asia 1. *Transbound Emerg Dis*, 67 (6): 3005-3015, 2020. DOI: 10.1111/tbed.13677
- Arju S, Nahar S, Ara MR, Sardar SA, Haque ME, Hossain MF, Basak P, Hossain MZ: Determination of TCID50 of O, A, Asia-1 serotypes foot-and-

- mouth disease virus to the production of trivalent vaccine in bioreactor. *Asian J Med Biol Res*, 8 (2): 63-68, 2022. DOI: 10.3329/ajmbr.v8i2.59507
28. **Habtewold WT, Welde N, Kenubih A, Getahun YA, Abayneh T, Getachew B, Woldemedhin W, Tesfaye Y, Tesfaw L, Getachew Y:** Effects of binary ethyleneimine and formaldehyde inactivation methods on foot-and-mouth disease virus vaccine immune responses and kinetics. *Sci Rep*, 15 (1):33788, 2025. DOI: 10.1038/s41598-025-01292-9
29. **Shojaee M, Zibaei S:** Purification of 146s antigen of foot-and-mouth disease (FMD) virus serotypes Aby using the sucrose gradient procedure. *Int J Infect*, 4 (1):e41322, 2016. DOI: 10.17795/iji-41322
30. **Abu-Elnaga H, Fawzy H, Farouk E, Ibrahim E, Gamil M, Zidan S:** Correlation between foot-and-mouth disease virus antigenic mass, titer and immune response in vaccinated sheep. *Benha Vet Med J*, 28 (2): 12-19, 2015. DOI: 10.21608/bvmj.2015.31855
31. **Bazid A-HI, El-Ashmawy WR, El-Sayed M:** Serum neutralization titers and protective efficacy induced by foot and mouth disease virus inactivated vaccine with different 146S particles concentrations in cattle. *Pak Vet J*, 36 (2): 204-208, 2016.
32. **Jamal SM, Bouma A, van den Broek J, Stegeman A, Chénard G, Dekker A:** Foot-and-mouth disease vaccine potency testing: The influence of serotype, type of adjuvant, valency, fractionation method, and virus culture on the dose-response curve in cattle. *Vaccine*, 26 (50): 6317-6321, 2008. DOI: 10.1016/j.vaccine.2008.09.021
33. **Zia MA, Dobson SJ, Rowlands DJ, Stonehouse NJ, Shah MS, Habib M:** Development of an ELISA to distinguish between foot-and-mouth disease virus infected and vaccinated animals utilising the viral non-structural protein 3ABC. *J Med Microbiol*, 71 (4):001516, 2022. DOI: 10.1099/jmm.0.001516